




INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 7 : C12N 15/29		A1	(11) International Publication Number: WO 00/32780
			(43) International Publication Date: 8 June 2000 (08.06.00)
(21) International Application Number: PCT/AU99/01079		[GB/AU]; 25A Bingham Circuit, Kaleen, ACT 2617 (AU). ROUSE, Dean, Thomas [AU/AU]; 131 Osburn Drive, MacGregor, ACT 2615 (AU).	
(22) International Filing Date: 2 December 1999 (02.12.99)		(74) Agent: GRIFFITH HACK; 509 St Kilda Road, Melbourne, VIC 3004 (AU).	
(30) Priority Data: PP 7469 3 December 1998 (03.12.98) AU 60/116,928 22 January 1999 (22.01.99) US		(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, VZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).	
(71) Applicant (for all designated States except US): COMMONWEALTH SCIENTIFIC AND INDUSTRIAL RESEARCH ORGANISATION [AU/AU]; Limestone Avenue, Campbell, ACT 2612 (AU).		Published With international search report.	
(71)(72) Applicant and Inventor: PEREZ, Pascual [FR/FR]; Laboratoire de Biologie Cellulaire et Moléculaire, Campus Universitaire des Cezeaux, 24, avenue de Landais, F-63107 Aubière (FR).			
(72) Inventors; and (75) Inventors/Applicants (for US only): BURN, Joanne, Elizabeth [AU/AU]; 84 A'Beckett Street, Watson, ACT 2602 (AU). PEACOCK, William, James [AU/AU]; 16 Brassey Street, Deakin, ACT 2600 (AU). DENNIS, Elizabeth, Salisbury [AU/AU]; 100 Hopetoun Circuit, Deakin, ACT 2600 (AU). SHELDON, Candice, Claire [AU/AU]; 1 Quartz Street, Sutton, NSW 2620 (AU). HELLIWELL, Christopher, Andrew			

(54) Title: CONTROL OF FLOWERING



(57) Abstract

The invention relates to control of flowering and reproduction in plants, and in particular to agents and methods for inducing or suppressing flowering. The invention provides isolated nucleic acid molecules which are useful for inducing flowering, particularly initiating early flowering, for delaying or suppressing flowering, or for manipulating the flowering period. In a first aspect, the invention provides an isolated nucleic acid molecule comprising a MADS box, which is capable of altering the flowering time of a plant. Preferably the nucleic acid molecule of the invention comprises a nucleotide sequence corresponding to a *FLOWERING LOCUS F (FLF)* gene. The nucleic acid molecule may be a genomic DNA, a cDNA, or a messenger RNA. The invention is applicable to any dicotyledonous or monocotyledonous plant species, including but not limited to decorative flower, vegetable, fruit, cereal, grass, tree, and other flowering species.

BEST AVAILABLE COPY

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Larvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

CONTROL OF FLOWERING

This invention relates to the control of flowering and reproduction in plants, and in particular to agents and methods for inducing or suppressing flowering. The invention provides isolated nucleic acid molecules which are useful for inducing flowering, particularly initiating early flowering, for delaying or suppressing flowering, or for manipulating the flowering period.

DETAILED DESCRIPTION OF THE INVENTION

The initiation of flowering in a plant occurs in response to internal signals, such as physiological age or levels of plant growth regulators, or may result from changes in environmental conditions such as day length or low temperature. It is well known that in a variety of plant species a crucial factor is day length, also known as photoperiod. In many plant species, including several ecotypes of the widely-used model plant species *Arabidopsis thaliana*, flowering is promoted by long day photoperiod, or by a period of low temperature (vernalization) (Napp-Zinn, 1985).

Control of flowering in both horticultural and crop plants represents a major problem in the agricultural industry, and is also a problem in forestry. Significant losses in yield of plants may result if non-uniform flowering of plants occurs; this applies both to field-grown and to glasshouse-grown plants. The problem is particularly acute for field-grown plants, which are frequently exposed to abnormal or unseasonal conditions which may result in induction of flowering at an inappropriate time. Efficient plant production requires the synchronization of flowering time between pollen donor and pollen receptor plants, and is particularly important to maximize market opportunities for glasshouse-grown plants.

Currently-available methods for regulation of flowering in plants are expensive and labour-intensive, and require the use of plant growth regulators, and/or controlled planting regimes and controlled-environment growth conditions. Consequently there is a need in the art for more efficient, cost-effective methods for controlling flowering time. These methods are applicable to a variety of commercially-significant plants species, including both horticultural plants, particularly those used in the cut-flower industry, and vegetable, cereal and other crop plants.

A class of genes known as MADS box genes encodes proteins which comprise a distinctive conserved DNA binding domain, known as the MADS box, which in certain cases has been demonstrated to bind to CC(A/T)₆GG DNA motifs. The MADS box genes encode a class of transcription factors, which was first identified in yeast and in mammals. Subsequently similar transcription factors were identified in a range of plants, including *Arabidopsis thaliana*, *Antirrhinum majus*, tomato, tobacco, petunia, corn, *Pinus* species and *Eucalyptus* species. In plants, the MADS box genes have a "K domain", which resembles the coiled-coil domains of keratin proteins, which are implicated in protein/protein interactions, an intervening (I) domain, and a carboxy terminal (C) domain. In plants the principal role of MADS box genes is in specifying inflorescence meristem identity, and floral organ identity and development. Certain MADS box genes have also been implicated as having roles in root and vegetative development.

We have now identified nucleic acid sequences comprising a MADS box in the model plant *Arabidopsis thaliana* which play a role in the control of flowering time. The effect on flowering depends on the degree of expression of the nucleic acid sequences.

SUMMARY OF THE INVENTION

In a first aspect, the invention provides an isolated nucleic acid molecule comprising a MADS box, which is capable of altering the flowering time of a plant.

5 In one preferred embodiment, the invention provides an isolated nucleic acid molecule which is capable of delaying the flowering of a plant. Preferably expression of the nucleic acid molecule in the plant, in the sense orientation under the control of a promoter
10 sequence, is capable of delaying the flowering of the plant.

In a second preferred embodiment, the isolated nucleic acid molecule of the invention is capable of accelerating the flowering of a plant. Preferably
15 expression of the nucleic acid molecule in the plant in the anti-sense orientation under the control of a promoter sequence is capable of accelerating the flowering of the plant.

Preferably the nucleic acid molecule of the invention comprises a nucleotide sequence corresponding to
20 a *FLOWERING LOCUS F* (FLF) gene. The nucleic acid molecule may be a genomic DNA, a cDNA, or a messenger RNA.

More preferably the nucleic acid molecule comprises the nucleotide sequence set out in any one of SEQ
25 ID NOS. 1, 2, 4, and 6 to 15, or a nucleic acid molecule capable of hybridizing thereto under at least low stringency hybridization conditions, or a nucleic acid molecule with at least 70% sequence identity to at least one of SEQ ID NOS. 1, 2, 4 and 6 to 15. Methods for
30 assessing ability to hybridize and % sequence identity are well known in the art. Even more preferably the nucleic acid molecule is capable of hybridizing thereto under high stringency conditions, or has at least 80%, most preferably at least 90% sequence identity. A nucleic acid molecule
35 having at least 70%, preferably at least 90%, more preferably at least 95% sequence identity to one or more of these sequences is also within the scope of the invention.

In a second aspect, the invention provides a vector comprising a nucleic acid molecule according to the invention. The vector may be a virus, bacteriophage, plasmid, or bacterium. In a particularly preferred embodiment, the vector is a T-DNA vector present in a bacterium of the genus *Agrobacterium*, in particular *Agrobacterium tumefaciens*.

In a third aspect, the invention provides a plant cell transformed with a nucleic acid of the invention.

In a fourth aspect, the invention provides a plant transformed with a nucleic acid molecule of the invention.

In a fifth aspect, the invention provides a method of isolating a nucleic acid molecule capable of altering the flowering time of a target plant, comprising the step of using a nucleic acid molecule of the invention, or a functional portion thereof, as a hybridisation probe or polymerase chain reaction (PCR) primer, and optionally detecting hybridisation. Suitable methods are very well known in the art. For example, we have demonstrated that the *Arabidopsis FLF* sequence described herein can be used to isolate the homologous sequence from *Brassica napus*.

In a sixth aspect, the invention provides an FLF polypeptide. Preferably the polypeptide is encoded by a nucleic acid molecule of the invention. More preferably the polypeptide has an amino acid sequence as set out in any one of SEQ ID NO: 3, 5, and 16 to 30, or has a sequence at least 70% identical thereto.

The polypeptide may be produced by expression of the FLF nucleic acid molecule in a convenient host, for example in a bacterial host such as *Escherichia coli*. Antibodies against the polypeptide, including monoclonal antibodies, may be produced using routine methods, and it will be clearly understood that antibodies to the FLF polypeptide are within the scope of the invention. Such antibodies are useful for screening plants for high or low levels of expression of FLF polypeptide. Suitable

screening methods including Western blotting and various forms of immunoassay, for example radioimmunoassay, ELISA, and chemiluminescent or fluorescent detection immunoassays.

Genes controlling developmental stages in plants, such as the gene associated with the nucleic acid of the invention, are highly conserved during evolution. Consequently the nucleic acid molecules and the methods of the invention are applicable to all plant species, whether the species is monocotyledonous or dicotyledonous. Thus the invention is generally applicable to flowering plants, including but not limited to ornamental, horticultural, agricultural and tree species. Methods for introducing exogenous DNA into plants of all these types, and for *in vitro* culture of plant tissue and regeneration of plant cells or tissues into whole plants, are known in the art. Methods for further generation and selection of commercially useful cultivars are also well known. Depending on the type of plant, it may be desirable to accelerate flowering *ie.* to induce early flowering, to synchronise flowering, to delay flowering or to suppress flowering.

For example it is desirable to suppress or delay flowering in many vegetable plants, in pasture grasses such as rye grass, or in sugar cane. Acceleration of flowering by induction of early flowering is desirable in a number of crop species, such as cotton, and in horticultural species.

We have surprisingly found that flowering can be delayed in proportion to the degree of expression of the nucleic acid molecule of the invention, and that early flowering can be induced by reducing the expression of this nucleic acid molecule.

Thus in a sixth aspect the invention provides a method of delaying flowering in a plant, comprising the step of introducing a nucleic acid molecule of the invention into cells of the plant, optionally such that expression of the nucleic acid molecule is under the control of an inducible promoter, and over-expressing the

nucleic acid molecule. Preferably the promoter is a tissue-specific promoter.

Preferably flowering is delayed for at least five days, preferably for at least twenty days, and more preferably for at least thirty days beyond the normal flowering period. Most preferably flowering is delayed for at least forty to fifty days. In at least some species it may be possible to achieve complete suppression of flowering. It will be appreciated that this further provides a method of inducing sterility in a plant.

According to a seventh aspect, the invention provides a method of inducing early flowering in a plant, comprising the step of reducing the degree of expression in the plant of a nucleic acid molecule of the invention. The reduction may be effected by any convenient means, including but not limited to transformation of the target plant with an anti-sense nucleic acid sequence, post-transcriptional gene silencing, ribozyme cleavage, disruption of the nucleic acid sequence using a transposable element or transposon, or by a procedure such as vernalisation. The person skilled in the art will readily be able to select the most suitable procedure for the particular plant species in question. Optionally the method of the invention may be supplemented by other treatments, such as an exogenous gibberellin.

Preferably flowering is at least five days earlier than the normal flowering period, more preferably at least ten days, and most preferably at least fifteen days earlier than the normal flowering period.

We have found that the degree of expression of *FLF*, and hence the flowering time, can be altered by modifying the activity of genes known to affect flowering time, including but not limited to *FCA*, *FVE*, *FPA*, *LD*, *FLD*, and *VRN2*. Therefore in both the sixth and seventh aspects of the invention, a further means of modifying the degree of expression of *FLF* is provided by modifying the activity

- 7 -

of one or more additional genes which affects flowering time, or vernalisation.

According to an eighth aspect, the invention provides a method of modifying the vegetative and/or floral phenotype of a plant, comprising the step of increasing the
5 level of expression of an *FLF* gene, thereby to modify the level of production or activity of a gibberellin in the plant.

Preferably the vegetative or floral phenotypic
10 characteristic is one which is regulated by gibberellic acid production or activity. More preferably the characteristic is related to plant architecture or fertility. For example, modification of gibberellic acid production and/or activity using the method of this aspect
15 of the invention may be used to produce dwarf or sterile plants. In one particularly preferred embodiment, the invention provides sterile plants. In a second preferred embodiment, the invention provides dwarf plants; more preferably the plant is a wheat plant.

20 In a number of embodiments of the invention, the nucleic acid molecule of the invention is operably linked to a promoter sequence capable of regulating the expression of the nucleic acid molecule; more preferably the promoter sequence is adapted to regulate expression in a eukaryotic
25 cell, most preferably a plant cell. The nucleic acid molecule of the invention may also be operably linked to a transcriptional terminator sequence.

Suitable promoter sequences are well known in the art, and include but are not limited to the CaMV 35S
30 promoter, a NOS promoter, the octopine synthetase (OCS) promoter, a subclover stunt virus promoter and the *Arabidopsis thaliana* ubiquitin gene promoter. The person skilled in the art will readily be able to select the most suitable promoter for a given purpose. In particular,
35 for some purposes an inducible promoter may be desirable, and these are also well known in the art. Suitable transcriptional terminator sequences active in plant cells

are also well known, and may be of bacterial, fungal, viral, animal or plant origin.

Suitable transcriptional terminators particularly suitable for use in the present invention include the
5 nopaline synthase (NOS) gene transcriptional terminator of *Agrobacterium tumefaciens*, the transcriptional terminator of the Cauliflower mosaic virus (CaMV) 35S gene, the zein gene transcriptional terminator from *Zea mays*, and the Rubisco small subunit (SSU) gene transcriptional terminator
10 sequences or subclover stunt virus (SCSV) gene sequence transcriptional terminators.

The nucleic acid molecule of the invention may be introduced into a plant cell or tissue by any suitable means. A variety of methods for introducing exogenous DNA
15 into plant tissue (transformation) are known. These include, but are not limited to, direct DNA uptake into protoplasts (Krens et al, 1982; Paszkowski et al, 1984), polyethyleneglycol-mediated uptake to protoplasts (Armstrong et al, 1990), electrophoresis (Fromm et al,
20 1985), microinjection of DNA (Crossway et al, 1986), microparticle bombardment of tissue explants or cells (Christou et al, 1988; Sanford, 1993), or T-DNA-mediated transfer from *Agrobacterium* to the plant tissue. Representative T-DNA vector systems are described in the
25 following references: An et al (1985); Herrera-Estrella et al (1983a, b); Herrera-Estrella et al (1985). These transformation methods are applicable to plant tissue culture, or may be employed with whole plants (*in planta* transformation). Again a person skilled in the art will be
30 able to select the most suitable method for any given plant.

Any plant tissue capable of subsequent clonal propagation, whether by organogenesis or embryogenesis, may be transformed with a vector of the present invention. The
35 particular tissue chosen will vary, depending on the clonal propagation systems which are most suitable for the species being transformed. Suitable tissue targets include whole

plant, leaf discs, pollen, embryos, cotyledons, hypocotyls, megagametophytes, callus tissue, existing meristematic tissue (eg. apical meristem, axillary buds, and root meristems), and induced meristem tissue (eg. cotyledon
5 meristem and hypocotyl meristem).

The vector of the invention may additionally comprise a dominant selectable marker to facilitate cell selection and plant breeding. A variety of suitable markers is known in the art, including but not limited to
10 the *NPTII* gene, genes encoding resistance to an antibiotic such as hygromycin or ampicillin or to a herbicide such as phosphinothricin or glyphosate; a gene encoding a polypeptide which confers stress tolerance, such as superoxide dismutase; or a visually-detectable marker, such
15 as green fluorescent protein or β -glucuronidase. The person skilled in the art will readily be able to select the most suitable marker for use in a specific case.

The invention is applicable to any dicotyledonous or monocotyledonous plant species, including but not
20 limited to decorative flower, vegetable, fruit, cereal, grass, tree, and other flowering species. Preferably the plant is selected from the group consisting of chrysanthemum, rose, gerbera, carnation, tulip, legumes such as soya bean, sugar beet, lettuce, cotton, oil seed
25 rape, coriander, *Lolium*, wheat, barley, maize, rice, pasture grasses, *Phalaris*, Canola and other Brassica species, *Linola* species, sugar cane, *Eucalyptus* species, pine and poplar. Forest species are to be understood to be within the scope of the invention.

30 For the purposes of this specification it will be clearly understood that the word "comprising" means "including but not limited to", and that the word "comprises" has a corresponding meaning.

The term "flowering time" as used herein means
35 the time at which floral meristem tissue is first visually detectable in the plant, for example by light microscopy or using the naked eye. The measured flowering-time includes

the time taken for the occurrence of the cellular processes in the differentiation of a floral meristem and subsequent cell divisions which enable such visual means to be used. The term "flowering time" also includes the time taken for the transition from a vegetative meristem to a floral meristem to occur, as measured visually, following the induction of flowering in the plant by the application thereto of a specific chemical, physical or environmental stimulus, such as a plant growth regulator, photoperiod or temperature regime, including the vernalisation of the plant. Alternatively flowering may be induced in response to an internal development signal in the plant. Those skilled in the art will be aware of the specific nature of such chemical, physical or environmental stimuli or internal developmental signals.

"Altering the flowering time" means that the time period in which floral meristem tissue is first visually detected in a plant is increased, decreased, or otherwise modified or regulated. Thus, flowering may be delayed, accelerated, inhibited, suppressed, or synchronized.

The term "meristem" refers to plant tissue in which cells are undergoing, or are capable of undergoing, rapid mitotic division followed by differentiation into cell types which are capable of forming a primordium which develops into an organ such as a leaf, root, stem, floral bud or other plant organ.

"Vegetative meristem" refers to a meristem in which the differentiation process produces a cell type which develops into a vegetative organ or non-reproductive organ, such as a leaf, petiole, bract, stem or root.

"Floral meristem" refers to a meristem in which the differentiation process produces a cell type which develops into an inflorescence meristem, a secondary inflorescence meristem, a floral organ or sexual reproductive organ, in which the meristem or organ, when developed, may comprise both reproductive and non-reproductive tissues, including, but not limited to,

anthers, stamens, stigmas, ovules, carpels, petals and sepals. "Bolt" refers to an inflorescence stem of a rosette plant, and "bolting" is the development of such a stem.

5 The term "derived from" means that a particular integer or group of integers has originated from a particular organism or species as specified herein, but has not necessarily been obtained directly from that source.

Representative low and high stringency conditions
10 of hybridisation as referred to herein are as follows:

High stringency: hybridization at 42°C in 50% formamide, 3 x SSC, 0.1% SDS, 20 x Denhardt's, 50 µg/ml salmon sperm DNA overnight and washed with a final wash of 0.1 x SSC, 0.1% SDS at 42°C.

15 Low stringency: hybridization at 28°C in 50% formamide, 3 x SSC, 0.1% SDS, 20 x Denhardt's, 50 µg/ml salmon sperm DNA overnight and washed with a final wash of 0.1 x SSC, 0.1% SDS at room temperature.

A "homologue" of a nucleotide sequence refers to
20 an isolated nucleic acid molecule which is substantially the same as the nucleic acid molecule of the present invention or its complementary nucleotide sequence, despite the occurrence within the sequence of one or more nucleotide substitutions, insertions, deletions, or
25 rearrangements.

An "analogue" of a nucleotide sequence means an isolated nucleic acid molecule which is substantially the same as a nucleic acid molecule of the present invention or its complementary nucleic acid, despite the occurrence of
30 any non-nucleotide constituents not normally present in the isolated nucleic acid molecule, for example carbohydrates, radiochemicals including radionucleotides, reporter molecules including, but not limited to digoxigenin, alkaline phosphatase or horseradish peroxidase.

35 A "derivative" of a nucleotide sequence means any isolated nucleic acid molecule which contains significant sequence similarity to the molecule or a part thereof. The

person skilled in the art will appreciate that the nucleotide sequence of the present invention may be subjected to mutagenesis to produce one or more single or multiple nucleotide substitutions, deletions and/or
5 insertions. Nucleotide insertional derivatives of the nucleotide sequence of the present invention include 5' and 3' terminal fusions, as well as intra-sequence insertions of single or multiple nucleotides or nucleotide analogues. Insertional nucleotide sequence variants are those in which
10 one or more nucleotides or nucleotide analogues are introduced into a predetermined site in the nucleotide sequence of the sequence, although random insertion is also possible; suitable screening of the resulting product is performed. Deletional variants are characterised by the
15 removal of one or more nucleotides from the nucleotide sequence. Substitutional nucleotide variants are those in which at least one nucleotide in the sequence has been removed, and a different nucleotide analogue inserted in its place.

20 Reference in this specification to a "gene" is to be understood in its broadest context, and includes:

(i) a classical genomic sequence comprising transcriptional and/or translational regulatory sequences and/or a coding region and/or non-translated sequences (ie.
25 introns and 5'- and 3'-untranslated sequences);

(ii) mRNA or cDNA corresponding to the coding regions (ie. exons), optionally additionally comprising 5'- or 3'-untranslated sequences of the gene; or

(iii) an amplified DNA fragment or other
30 recombinant nucleic acid molecule produced *in vitro*, and comprising all or a part of the coding region and/or 5' or 3'-untranslated sequences of the gene.

The term "gene" is also used to describe synthetic or fusion molecules encoding all or part of a
35 functional product. A functional product is one which comprises a sequence of nucleotides or is complementary to a sequence of nucleotides which encodes a functional

polypeptide, in particular the *FLF* polypeptide of the invention or a homologue, analogue or derivative thereof.

In some of the examples herein the *FLF* gene is referred to as gene B. These two terms are synonymous.

5

BRIEF DESCRIPTION OF THE FIGURES

Figure 1 is a photograph showing wild-type C24 (left), the late flowering T-DNA tagged *flf* mutant (middle) at 70 days after germination, and *flf* mutant at 150 days, showing the domed shape caused by vegetative bolts (right). The bar represents 5 cm.

Figure 2 shows segregation of two T-DNA inserts with the late flowering phenotype

A. Genomic DNA isolated from a T₂ population segregating for early (E), late (L) and very late (VL) flowering, digested with *EcoRI* and probed with the *NPTII* gene.

B. Physical map of the two T-DNA inserts linked to the *FLF* locus, showing their orientation. *EcoRI* sites are labelled RI; LB and RB represent the left and right borders, respectively, of the T-DNA. The triangle symbol represents the site of deletion of 30 bp to the right of the T-DNA. The arrows represent the direction of transcription of the genes.

C. Representation of a 27 kb region of *Arabidopsis* mutant DNA containing the gene A and gene B loci, showing the location of the T-DNA inserts. DNA fragments from the flanking plant DNA (probes 2 and 3) were used as probes to isolate cDNA clones. *HindIII* (H), *BamHI* (B) and *EcoRI* (R1) restriction enzyme sites are indicated. The positions of gene A and gene B are shown, and their directions of transcription are indicated by arrows.

D. The 6.5kb and 6.8 kb *BamHI* fragments, isolated from a genomic library of wild-type C24 with probes 1 and 2 from Figure 2C, spanning the site of the T-DNA insertions. Restriction sites are as in Figure 2C.

35

Figure 3 shows levels of expression of gene A and gene B in 30 day old wild-type C24 plants (lane 1), hemizygous (lane 2) and homozygous (lane 3) *flf* mutant plants.

5 Figure 4 is a photograph showing 35S::*FLF* T1 transgenic plants in C24 (left) and Landsberg *erecta* (right) ecotypes. The C24 transgenics were either early-flowering (back) or late-flowering (front). The Landsberg *erecta* transgenics were either late-flowering (left) or
10 flowered at normal time (right).

Figure 5 shows the *FLF* gene structure and the expression pattern of the *FLF* transcript

A. Genomic structure of the *FLF* gene, showing location and size of introns and location of the MADS box, intervening domain (I), K domain (K) and carboxy terminal
15 domain (C). The numbers below the line represent the number of base pairs in each exon.

B. Pattern of expression of *FLF* mRNA in C24 plants: roots (R) and rosette leaves (RL) from *in vitro*
20 grown vegetative plants, cauline leaves (CL), bolt stems (BS), floral apex and buds (B) from soil grown plants with bolt stems between 1 and 5 cm. Mature flowers (F) and siliques (S) were collected from older plants. Plants were grown under 16 h photoperiod conditions. The RNA gel blots
25 for B-F were probed with a riboprobe transcribed from the *FLF* (Gene B) cDNA clone linearised to remove the MADS box region. The ethidium bromide-stained ribosomal bands are shown as a loading control in B-F.

C. Expression level of *FLF* mRNA in whole C24
30 or *flf* mutant plants, harvested every 10 days (as indicated by the numerals) until the majority of the C24 plants were bolting (50 days under these growth conditions).

D. Expression of *FLF* mRNA in C24 (lane 1-6) and *flf* (lane 7-12) plants grown for 21 days in 8 h
35 fluorescent photoperiod, and then at the end of the 21st photoperiod either kept in the same conditions (SD; lane 1, 2, 7, 8) or transferred to continuous dark (CD; lanes 3, 4,

- 15 -

9, 10) or continuous light (CL; lanes 5, 6, 11, 12). Plants were harvested either just prior to what would have been the start of the following photoperiod (dawn; lane 1, 3, 5, 7, 9, 11), or just prior to the end of the photoperiod (dusk; lanes 2, 4, 6, 8, 10, 12). Transcript levels were a little higher at the start of the photoperiod, but this pattern was not altered in the mutant.

E. Effect of gibberellic acid (GA_3) treatment and vernalization on the *FLF* transcript in C24 (lanes 1-3) and *flf* mutant (lanes 4-6) seedlings. RNA was isolated from 12 day old seedlings that had either had no treatment (C; lanes 1 and 4), been grown on medium containing 10^{-5} M GA_3 (G; lanes 2 and 5) or had a pretreatment of 3 weeks at 4°C (V; lanes 3 and 6).

F. *FLF* expression in rosette leaves of C24 and the early-flowering antisense methyltransferase line 10.5 (T3 generation) harvested soon after bolting.

Figure 6 shows

A. Genomic DNA isolated from individual *flf*, *efSL3* (M2), *efSL4* (M2) and C24 plants digested with *EcoRI* and probed with a probe directed to the 3' region of *Ac*. The DNA for the *flf* sample was extracted from plants which contained a third T-DNA band, hence the band at about 8 kb. The presence of this third band had no effect on flowering time, and is therefore irrelevant.

B. As for **A**, except that the probe was probe 4 (see Figure 2C). The DNA for the *efSL3* (M2) and *efSL4* (M2) samples was extracted from bulked M2 plants which contained neo-lates as well as early-flowering mutants. Therefore there is some of the 2.7 kb band present in these DNA extracts. Other DNA isolated from individual early-flowering plants does not contain a band at 2.7 kb.

C. Location of *Ac* insertion in intron I. The nucleotide positions are given below, taking the A of the ATG as nucleotide 1.

D. Expression level of *FLF* gene in 15 day old rosette leaves of C24 (lane 1), *flf* (lane 2), *ef* SL3 (lane 3), *ef* SL4 (lane 4). The M2 early flowering mutants had just started to bolt, whereas the other plants remained vegetative. The ethidium bromide-stained ribosomal bands are shown as a loading control.

Figure 7 shows gel blots from neo-late plants.

A. Genomic DNA was isolated from the 6 neo-late mutant plants and from *ef*SL3 (M2), *ef*SL4 (M2), *flf* and C24 plants, and digested with *Eco*RI. The DNA gel blot was probed with the 3' region of *Ac*.

B. Total RNA was isolated from a mixture of rosette and cauline leaves from the 6 neo-late plants, *flf* and C24. The RNA gel blot was probed as in Figure 5. The ethidium bromide-stained ribosomal bands are shown as a loading control.

Figure 8 shows expression of the *FLF* gene in ecotypes and late-flowering mutants.

Total RNA was extracted from 12 d old seedlings and RNA gel blots were probed as in Figure 5. The ethidium bromide-stained ribosomal bands are shown as a loading control in A-C.

A. Expression in a number of different *Arabidopsis* ecotypes

B. Expression in Landsberg *erecta* (L.*er.*) and Landsberg *erecta* lines which contain late alleles at either the *FRI* (L.*er.*-*FRI*^{Sf2}) or *FLC* (L.*er.*-*FLC*^{Sf2}, L.*er.*-*FLC*^{Col}) loci.

C. Expression in late-flowering mutants in either L.*er.* ecotype (*fca*, *fve*, *fpa*, *gi*, *co*, *fha*, *fwa*, *fd*, *fe*, *ft*) or *Ws* (*ld*). The mutants *vrn1* and *vrn2* were isolated in the *fca* background, and only *vrn1* has been segregated away from the *fca* mutant locus.

Figure 9 shows the partial genomic sequence of an *FLF*-like sequence from *Brassica napus*, showing the location

of exons, and the predicted sequence of the translated product.

Figure 10 shows a comparison of the predicted translated product of a *Brassica napus* *FLF*-like sequence (top lines), and the predicted *FLF* translation product from *Arabidopsis thaliana*, showing identical amino acids (|), highly conserved amino acids (:), and conserved amino acids (.) .

10 DETAILED DESCRIPTION OF THE INVENTION

The invention will now be described in detail by way of reference only to the following non-limiting examples, and to the figures.

15 GENERAL METHODS

Plant Material and Growth Conditions

Arabidopsis was grown either in pots containing a mixture of 50% sand and 50% loam, or aseptically in test-tubes or petri dishes containing a modified Murashige and Skoog (MS) medium (Langridge, 1957). Unless otherwise stated, all plants were grown in artificially lit cabinets at 21° or 23°C, under long day (16 hr light, 8 hr dark) conditions using cool white fluorescent lights at an intensity of 200 $\mu\text{M m}^{-2}\text{s}^{-1}$.

Plants were vernalized by germinating seed in the dark for either 3, 4 or 8 weeks at 4°C. Following this cold treatment, seedlings were transferred to long day photoperiods at 23°C and times to flower, measured as the time until stem elongation (bolting) was observed, were determined, beginning from the first day at the higher temperature.

Arabidopsis Transformation

Arabidopsis was transformed either by root transformation (Valvelkens et al., 1988) for the generation of the *flf* mutant and gene A transgenic plants, or by in planta transformation (Bechtold et al, 1993) for the gene B

transgenic plants. The late-flowering mutant (*flf*) arose during transformation of the early-flowering ecotype C24 with a modified binary vector pBinΔAc (Finnegan et al, 1993). This vector contains the neomycin

- 5 phosphotransferase II (*NPTII*) gene under the control of the nopaline synthase (*NOS*) promoter, together with a deleted maize Ac transposable element inserted within the untranslated leader of the β -glucuronidase gene, in the reverse orientation to the direction of transcription.

10

Example 1 Identification of a Mutant Locus Associated With a Late Flowering Phenotype

- Following transformation of the *Arabidopsis thaliana* ecotype C24 with a T-DNA construct containing an
15 Ac transposable element, individual T₀ plants, resistant to kanamycin, were allowed to self-pollinate, and the T₁ progeny screened for families that flowered significantly later than parental C24 plants.

- Some of the plants of family 14-58 flowered after
20 70 days compared to 30 days for the C24 control plants. Segregation analysis of the progeny of a selfed late-flowering T₁ plant from family 14-58, showed 53 "late" (flowering time >70 days) compared to 15 "early" flowering plants (flowering time 30 days). The result fits a 3:1
25 segregation ratio ($\chi^2=0.313$ P>0.5), and is consistent with the late-flowering phenotype being a consequence of a single mutation. We have designated the mutant locus *FLOWERING LOCUS F (FLF)*.

- Within the segregating progeny, the late-
30 flowering plants could be further differentiated into two classes; "lates", flowering between 70 to 90 days and "very lates", which flowered later than 150 days. Some of the "very late" plants had not flowered after a year of growth. The "very late" *flf* mutant, shown in Figure 1, produced
35 leaves at a rate similar to non-transformed C24, and had many more leaves at flowering than C24. After several months of growth, bolts arose from the internodes between

- 19 -

rosette leaves. These bolts elongated approximately two to three cm and formed aerial rosette structures, giving the mutant plants a dome-like appearance, similar to that described for the *fld* mutant (Chou and Yang, 1998).

5 The late-flowering phenotype observed is more extreme than any of the previously-reported late-flowering mutants and ecotypes (Koornneef et al, 1991). Progeny tests showed that selfed "very lates" produced only "very late" progeny, whereas selfed "late" plants segregated 1:2:1 for
10 very late, late and early flowering plants. This segregation pattern is consistent with a semi-dominant mutation, with the lateness in flowering being proportional to the level of gene product.

15 Example 2 Construction and Screening of Genomic Libraries

A genomic library of the *flf* mutant was constructed by partial digestion of total plant DNA with the restriction enzyme *Sau3AI* and ligation into the phage
20 vector λ EMBL4. The resulting library was screened using a 32 P-dCTP labelled probe of the *NPTII* gene (Feinberg and Vogelstein, 1983). Four positive phage clones were purified and restriction mapped. Together these spanned 27kb of plant DNA flanking the site of insertion of T-DNA.
25 A 2.3kb *Bam*HI-*Eco*RI and a 2.7 kb *Eco*RI fragment (probes 1 and 2 respectively, Figure 2C) isolated from this flanking plant DNA were subsequently used to probe a genomic library of wild-type *Arabidopsis* C24, made from *Bam*HI-digested DNA and cloned into λ EMBL4. Probe 1 hybridized to a genomic
30 clone containing 6.5kb of plant DNA spanning the T-DNA insertion site, and probe 2 hybridized to a genomic clone containing 6.8 kb of adjacent sequence.

Example 3 Isolation of the *FLF* Gene

35 Two T-DNAs segregate with the very late flowering phenotype
The very late-flowering phenotype segregates with two T-DNA bands identifiable by Southern analysis, which

- 20 -

were designated bands 1 and 5. Southern blotting showed that bands 1 and 5 are inverted and adjacent:

RB ←—— LB LB →—— RB
 5 1 5
 (LB, left border; RB, right border)

The size of the bands, combined with sequence analysis, places the smaller band (band 5) closest to the *FLF* gene. Recombinant inbred lines were used to map the *FLF* region to the top of chromosome 5, 4 cM from RFLP marker 447. This places the *FLF* gene near *FLC*, a gene known to control flowering time in ecotypes of *Arabidopsis*.

A 27 kb segment of genomic DNA from the mutant around the site of T-DNA insertion was mapped. C24 genomic clones covering 13 kb around the T-DNA insertion site in the mutant were sequenced. Two probes, on either side of the T-DNA insert, were used to screen cDNA libraries. A 4.6 kb EcoRI/BamHI fragment from a C24 genomic clone (containing 4.4kb of sequence "upstream" of the T-DNA insertion site and 0.2 kb "downstream" of the T-DNA insertion site) was used to isolate cDNA clones identifying "gene A" as a transcribed region. A 2.7 kb EcoRI fragment, covering the region 0.4 kb "downstream" to 3.1 kb downstream of the T-DNA inserts, was used to isolate cDNA clones identifying "gene B" as a transcribed region.

Comparison of part of the intergenic region sequence between ecotypes C24 and Ws revealed an insertion into the C24 DNA of an approximately 200 bp sequence, 420 bp to the "right" of the stop codon of gene A and 120 bp to the "left" of the T-DNA insertion. The sequence is present in both C24 wild-type and the *flf* mutant. The 200 bp insertion shows 100 % homology to ORF167 of the *Arabidopsis* mitochondrial genome. By PCR analysis we have determined that this sequence is absent in Landsberg *erecta* and Columbia ecotypes. The significance of this inserted DNA segment is unknown.

A map of the overall region is shown in Figure 2B.

Example 4 cDNA Libraries

5 In order to identify expressed genes closely
linked to the T-DNA, three *Arabidopsis* cDNA libraries
(Elledge et al, 1991; Weigel et al, 1992; Newman et al,
1994) were screened, using probes to plant DNA around the
T-DNA insertion site (probes 2 and 3, Figure 2C). Two
10 classes of cDNA clones were isolated. These were
respectively designated gene A and gene B. Gene B was
subsequently re-designated as *FLF*. Two gene A cDNAs were
isolated with a 4.6 kb EcoRI-BamHI fragment (probe 3,
Figure 2C) from a screen of 200,000 λ Yes clones; however,
15 no clones were isolated with the 2.7 Kb EcoRI fragment
(probe 2, Figure 2C). The gene A cDNAs were subcloned from
the phage by site-specific recombination, using the CRE
protein (provided by the *E. coli* strain BNN132) and the lox
sites within the vector (Elledge et al, 1991). The larger,
20 almost full-length gene A cDNA was further subcloned into
pBluescript SK(-) (Stratagene). Full-length gene A cDNAs
were subsequently isolated by screening a *Landsberg erecta*
flower cDNA library (Weigel et al, 1992). The mutant and
wild-type genomic clones corresponding to the isolated cDNA
25 were also subcloned as smaller fragments into pBluescript
SK(-). As no cDNA clones were isolated with probe 2
(Figure 2C) from either cDNA library, a third library was
screened. Four full-length gene B cDNA clones were
isolated from a λ PRL2 cDNA library derived from different
30 tissues and developmental stages (Newman et al, 1994). All
cDNAs and the mutant and wild-type genomic clones were
sequenced on both strands by the dideoxy chain termination
method (Sanger et al, 1977) using fluorescent primers
(Brumbaugh et al, 1988). The University of Wisconsin GCG
35 software package was employed for sequence analysis
(Devereux et al, 1984). The nucleotide and predicted

protein sequences were used to search the GenBank database for any homologous sequences; none were found.

Example 5 Construction of 35S::gene A plasmid

5 As the larger of the initially isolated gene A cDNA clones lacked the AT of the ATG of the start codon, oligonucleotide-directed mutagenesis was employed to generate a 200bp fragment from the 5' end of the cDNA which contained the absent nucleotides. Two oligonucleotides
10 were synthesized on an Applied Biosystems DNA Synthesizer for this purpose:-

- (1) 5' AAGCCGCGGACAATGGAAGCTGTAAGATGC 3'
(2) 5' GAGAGGCTG**GTTA**ACCGGAG 3'.

15 The nucleotides indicated in bold show the locations of *SacII* and *HpaI* restriction sites within the primers. The amplification reaction was carried out in a 10 µl final volume that contained 2 µM of each
20 oligonucleotide primer, 200pg of *HindIII*-cleaved cDNA as a template, 0.2 units of Taq polymerase and 125 µM of each of the four deoxynucleotides. Conditions for the amplification were as follows: 95°C for 2 mins, 5 cycles consisting of 15 s denaturation 95°C, annealing at 40°C for
25 30 s, and polymerization at 72°C for 1 min, followed by 25 cycles where the annealing temperature was raised to 50°C for 15 sec and finally 30°C for 1 min. The resulting
30 200 bp PCR fragment was cloned into *SacII* and *HpaI* sites in the original cDNA plasmid, and then sequenced to ensure
35 that no mutations had been introduced during the amplification procedure. Sense binary constructs were made by digesting the full length cDNA with *EcoRI* and *SacII*, end filling the recessed termini using the Klenow fragment of DNA polymerase I, and ligating the released 1.4 kb fragment into the *SmaI* site of the expression vector pDH51 (Pietrzak et al, 1986). This places the expression of the *FLF* cDNA under the control of the CaMV 35S promoter. Recombinant

- 23 -

plasmids, containing the cDNA in the desired orientation, were cleaved with *EcoRI* and cloned between the right and left border sequences of the binary vector pBin19 (Bevan, 1984). The binary construct was transferred to

5 *Agrobacterium tumefaciens* strain AGL1 (Lazo et al, 1991) by triparental mating, employing pRK2013 as the helper plasmid. Roots of wild-type C24 plants were transformed (Valvelkens et al, 1988) using the *NPTII* gene as a selectable marker to identify transgenic plants.

10

Example 6 Construction of 35S::gene B plasmid

A binary construct containing gene B under the control of the CaMV 35S promoter was generated by cloning a *XhoI/SpeI* digested PCR product, amplified using the gene B

15 cDNA clone as template with the primers, using methods similar to those described in Example 5:

5' CCGCTCGAGCTTAGTATCTCCGGCG 3' and
5' GGACTAGTCGCCCTTATCAGCGGA 3',

20

in which restriction sites are shown in bold, and the sequence hybridizing to gene B cDNA is underlined, into *XhoI/SpeI* digested pART7 (Gleaves, 1992) containing the CaMV 35S promoter. The 35S::gene B cassette was then

25 subcloned using *NotI* into pART27 (Gleaves, 1992) and introduced into *A. tumefaciens* strain GV3101 (Koncz and Schell, 1986) as described above. Transgenic plants were generated by *in planta* transformation (Bechthold et al, 1993)

30

Example 7 DNA Gel Blot Analysis

Total genomic DNA was isolated either by following the cetyltrimethylammonium bromide (CTAB) procedure (Dean et al, 1992), or as described by McNellis

35 et al. (1998). 2-3 µg of DNA was digested with the appropriate restriction enzyme, electrophoresed on an 0.8% agarose gel, and blotted onto Hybond N+ membranes

(Southern, 1975). Probes were labelled with ^{32}P -dCTP using the random primer method (Feinberg and Vogelstein, 1983).

The NPTII probe was generated as described above. The 3'Ac probe was a SphI fragment (Lawrence et al, 1993), and probe 4 was generated by amplification of the wild-type genomic clone with primers:

5'-GTATAGGGCACATGCCC-3' and
5'-CACTCGGAGCTGTGCC-3'.

This results in a 570 bp subset of probe 2 sequence, lacking the MADS box, to eliminate cross-hybridization.

15 Example 8 RNA Extraction and RNA Gel Blot Analysis

Total RNA was extracted from approximately 1g of plant tissue, following the method of Longemann et al. (1987). 10-20 μg of total RNA was electrophoresed on 2.2 M formaldehyde/agarose gels, and blotted onto Hybond N nylon filters. T7 or SP6 polymerase transcription of the linearised gene A or gene B plasmid template (containing the complete cDNA for gene A, linearised to remove MADS box for gene B) was used to generate antisense ^{32}P -dUTP labelled riboprobes. Filters were hybridized as described by Dolferus et al (1994), washed with a final solution of 0.1xSSC, 0.1% SDS at 65°C. For gene A Northern blots it was necessary to treat filters with RNase A as previously described (Dolferus et al, 1994) to avoid ribosomal trapping. The filters were exposed to phosphor screens for quantification of signal intensity using a phosphorimager (Molecular Dynamics, USA). RNA size markers were used to determine the size of the gene A and gene B transcripts.

35 Example 9 RFLP Mapping

DNA from sixty-four recombinant inbred lines (Lister and Dean, 1993) was digested with *Bam*HI, and Southern blots were probed with gene A. The Mapmaker

program was employed to compare the data with RFLP data for 68 mapped markers. Fine mapping of gene A was performed using DNA from F₂ plants generated from a cross between Landsberg erecta and the *flf* mutant. A *Hpa*I digest of 62 F₂ DNA was probed with the chromosome 5 RFLP marker 447 (Chang *et al.*, 1988). A restriction fragment length polymorphism (RFLP) was found between the parental lines, Landsberg erecta and Columbia, by digestion of genomic DNA with *Bam*HI and probing with the gene A cDNA.

Example 10 Insertion Of Two Inverted and Adjacent T-DNAs
Produces A Partial-Dominant Late Flowering
Mutant

DNA gel blot analysis of a late flowering T₁ parent showed five T-DNA inserts. Figure 2A shows a DNA gel blot of 16 segregating progeny plants of a selfed, hemizygous mutant plant derived from this T₁ parent. In a total of 70 progeny, only two inserts segregated with the late-flowering phenotype (bands 1 and 5); "late" plants were hemizygous for these two T-DNAs, and "very late" plants were homozygous for the same two inserts. The two linked T-DNAs were segregated away from the other T-DNA inserts by backcrossing to non-transformed C24. Plants containing only the two T-DNA inserts (Figure 2A, bands 1 and 5) were identified by DNA gel blot analysis, and further analysis showed that the two inserts were adjacent and in inverted orientation (Figure 2B). DNA gel blot analysis, using probes derived from segments of the T-DNA construct and *Ac* demonstrated that no movement of the *Ac* transposable elements had occurred.

A genomic DNA library from a late flowering plant containing only the two linked T-DNA inserts was screened with an *NPTII* probe to isolate DNA segments spanning the site of T-DNA insertion. Three overlapping clones were isolated from the left side of the T-DNA. The longest of these, together with one clone isolated from the right

side, is depicted in Figure 2C. These clones accounted for a total of 27 kb of plant DNA spanning the site of the T-DNA insertion. C24 genomic clones were isolated from a genomic DNA library, which was prepared by digestion of total *Arabidopsis* DNA with the restriction enzyme *Bam*HI and ligation of the digest into the phage vector λ EMBL4, using probes 1 and 2 (Figure 2C). Clones containing a 6.5kb *Bam*HI fragment spanning the insertion site and a 6.8 kb *Bam*HI fragment downstream of the insertion site were characterized (Figure 2D). The genomic DNA sequence, cDNA sequence, and predicted protein sequence are set out in SEQ ID NO. 1, SEQ ID NO. 2 and SEQ ID NO. 3 respectively. The genomic DNA sequence includes about 2 kb of promoter sequence, and 6 introns.

The chromosomal location of the *FLF* region was determined using 64 F9 recombinant inbred lines and data for several known markers (Lister and Dean, 1993). The Mapmaker program located the *FLF* region to the top of chromosome 5, 4cM from the RFLP marker 447, placing the gene in the vicinity of three other genes, *TFL*, *FLC* and *EMF1*, which are involved in the regulation of flower initiation (Shannon and Meeks-Wagner, 1991; Lee et al, 1994b; Koornneef et al, 1994; Sung et al, 1992).

Example 11 Two Genes Flanking the T-DNA Inserts Have Increased Expression In The *flf* Mutant

Transcriptionally active regions in the plant DNA spanning the site of T-DNA insertion were identified by screening a cDNA library with probes derived from either side of the T-DNAs (probes 2 and 3, Figure 2C). Two overlapping partial-length cDNA clones were isolated with probe 3; full-length clones containing inserts of approximately 1.5 kb were isolated by screening a second cDNA library. Four identical full-length cDNA clones containing inserts of 1.0 kb were isolated with probe 2. Comparison of cDNA and genomic sequence revealed that the T-DNAs had inserted between the two transcribed regions,

591 bp downstream of the polyadenylation site of gene A and 2.3 kb upstream of the start codon of gene B. Comparison of the mutant and C24 genomic sequences revealed a 30 bp deletion immediately downstream of the insertion site, with no further differences identifiable.

As neither gene was disrupted by the insertion of the T-DNAs, we investigated whether the expression of the genes was altered in the mutant. Figure 3A shows an RNA gel blot performed using RNA isolated from 30 d old leaf tissue of ecotype C24, and hemi- and homozygous *flf* mutant plants grown under identical conditions. As shown in Figure 3 antisense riboprobes specific for either gene A or gene B revealed that both genes are more highly expressed in the *flf* homozygote leaf tissue than in ecotype C24, with the 1.5 kb gene A transcript being approximately 10 times more highly expressed in the mutant and the 1.0 kb gene B transcript being approximately two-fold overexpressed in the homozygous mutant. The hemizygous mutant has an intermediate level of expression of both genes. Without wishing to be limited by any proposed mechanism, we believe that insertion of the T-DNA complete with Ac elements has caused this over-expression.

Example 12 Transgenic Plants Over-Expressing Gene B Have Altered Flowering Time

In order to determine which gene is responsible for the late-flowering phenotype, we transformed C24 with constructs containing either gene under the control of the CaMV 35S promoter. 49 transgenic lines were generated with the gene A construct, and flowering time was assessed in the T2 generation. The majority of the transgenic lines showed no variation from wild-type flowering time; however, a few lines were slightly late-flowering. As shown in Table 1, four of the transgenic lines were significantly later flowering than the C24 wild-type. However, there was no correlation between time to flowering and the level of gene A expression. 23 transgenic lines were generated with

the gene B construct, and differences in flowering time compared to C24 were apparent in the T1 generation, ie. in the primary transformants. 17 T1 plants flowered earlier (range 15-25 d) than non-transformed C24 (30 d) under these conditions; eight of these showed either full or partial sterility. Four flowered at around the same time as C24, and two had not flowered after 90 d. Examination of the level of gene B mRNA transcript in kanamycin-resistant progeny of two early-flowering T1 plants, and rosette leaves from two late-flowering T1 plants, revealed a high level of gene B expression in all the transgenic plants.

Table 1
Flowering time of 35S::gene A transgenic plants

	Flowering time	Relative expression level of gene A mRNA
C24	29.9 \pm 0.6	1
B2	38.3 \pm 0.9	2
A53	35.8 \pm 0.8	20
A54	38.4 \pm 0.5	8
A93	35.4 \pm 0.5	4
<i>flf</i>	>>50	10

Transgenic and *flf* mutant seeds were germinated on MS plates + 50 μ g/ml kanamycin, and C24 seeds were germinated on MS plates. At least 20 12 d seedlings were transplanted into individual soil pots and grown at 23°C under fluorescent lights (16 h light, 8 h dark). Flowering time was recorded as the number of days to stem elongation. Total RNA was extracted from 14 d *in vitro* grown seedlings and used for quantitation of *FLF* expression levels.

Table 2

Flowering time of 35S::gene B T1 transgenic plants

	Flowering time	Relative expression level of gene B mRNA
C24	30 d	1
B4	18 d	> 10
B5	18 d	> 10
B11	> 80 d	> 10
B12	> 80 d	> 10
Landsberg <i>erecta</i>	20 d	n.d.
B36	> 80 d	> 10
B45	> 80 d	> 10
<i>flf</i>	> 80 d	2

- 5 Transformant seeds, harvested from *in planta* transformed plants, were selected on MS plates containing 50 µg/ml kanamycin. Kanamycin resistant T1 seedlings were transferred to soil at 20 d. The level of gene B transcript was determined in kanamycin resistant T2 plants
- 10 for B2 and B5, and from young leaves from the T1 late-flowering plants (B11, B12, B36, B45). n.d.: not detectable.

- 15 It is surprising that over-expression of gene B gave two completely opposite phenotypes. In order to clarify this, we generated transgenics containing gene B under the control of the 35S promoter in ecotype Landsberg *erecta*. Of the 24 T1 lines generated, none flowered earlier than wild-type Landsberg *erecta*, 12 had not bolted
- 20 after 70 days, and 3 bolted after about 40 days, compared with 25 days for Landsberg *erecta*. Two of the three lines which bolted after 40 days exhibited floral abnormalities and partial sterility. Total RNA was isolated from rosette

leaves of two non-flowering T1 plants (B36, B45; Table 1), both of which had high expression levels of the transgene.

Therefore in Landsberg *erecta* over-expression of gene B causes a delay in flowering time, whereas in C24 it causes either a delay in flowering, or causes the plants to flower significantly earlier. This may be mediated by a dominant negative effect or by a form of post-transcriptional gene silencing. Analysis of protein expression levels is being pursued in order to clarify this point. Presumably a difference in the genetic background of the two ecotypes is responsible for the difference observed between ecotypes.

These results demonstrate that over-expression of gene B causes late-flowering, whereas gene A has little effect on flowering time, indicating that over-expression of gene B is the most likely cause of the late-flowering *flf* phenotype and that this gene encodes a dosage-dependent repressor of flowering. Gene B will hereafter be referred to as *FLF*.

Example 13 Anti-Sense Constructs

Anti-sense plant constructs have been generated using an anti-sense *FLF* gene construct under the control of the CaMV 35S promoter. A 35S::*FLF* antisense binary construct was generated by cloning the EcoRI/SpeI digested PCR product amplified with primers

CGGAATTCTCACACGAATAAGGTAC and

GGACTAGTGGTCAAGATCCTTGATC

as described for the 35S::*FLF* construct. This amplified the region downstream of the MADS box, so that the antisense construct lacks the MADS box region. The PCR product was cloned into pART7 and pBART 27 (which is a derivative of pART27), and transgenic plants were generated as described above, except that the *Bar* gene was used as the selectable marker.

25 T1 C24 transgenic plants were generated with a construct in which the 3' end of the *FLF* gene, in the

- 31 -

antisense orientation, was under the control of a 35S promoter (35S::*FLFAS*). Approximately half of the T1 plants had flowered before 20 days of growth, compared to 30 days for the non-transformed strain

5 Transgenic plants were produced in the C24 and the Columbia ecotypes. Of the six T1 plants produced in the Columbia ecotype, three bolted earlier than wild-type Columbia. Wild-type C24 plants bolt at about 30 days, and wild-type Columbia plants bolt at about 20 days.

10 These results indicate that the antisense construct acts to decrease flowering time, presumably by decreasing the level of the *FLF* transcript.

RNA gel blot analysis of early-flowering plants from three T2 C24 antisense lines revealed that the level
15 of *FLF* transcript was considerably lower than in non-transformed C24, confirming that the antisense construct was acting to decrease flowering time by decreasing the level of *FLF* transcript.

20 Example 14 The *FLF* Gene Is A Novel MADS Box Gene

The *FLF* cDNA sequence has strong homology to a class of transcription factors known as MADS box genes. The *FLF* sequence shows greatest similarity to the MADS gene AGL14 in the M-I-K domain, but over the entire cDNA
25 sequence it shows greater similarity to *CAL* (*CAULIFLOWER*) and *AP1* (*APETALA1*). The location of the MADS box, I domain, K domain and C terminal domain are indicated in Figure 5A. The K domain is typical of those of other plant MADS genes. Comparison of the genomic sequence (SEQ ID
30 NO. 1) and cDNA sequence (SEQ ID NO. 2) of the *FLF* gene revealed the presence of 6 introns, with intron I being 3.5 kb. The predicted protein (SEQ ID NO. 3) is 196 amino acids long, which is shorter than the proteins encoded by most MADS box genes.

35 One of the main roles of MADS box genes in plant development which has been described to date is in specifying floral organ identity. Other roles for MADS-box

genes include specifying root architecture and vegetative growth. To investigate whether *FLF* also has other roles in addition to its role in controlling the time of flowering, we examined its expression in a range of tissues. Figure 5B confirms the high expression of the *FLF* gene in vegetative rosette leaves, and reveals a strong expression in roots and lower expression in floral tissues, suggesting possible further roles for the *FLF* gene. No root phenotype has been observed in the transgenic lines. However, a number of lines had reduced fertility, which appeared to be caused by a lack of pollen in C24 lines or by abnormal carpels in Landsberg erecta lines. However, as the early Ac plants (see later) did not show these phenotypes it is unclear whether this is caused by change in expression of the *FLF* gene.

The expression of the *FLF* gene is lower in post-vegetative tissues than it is in vegetative rosette leaves. We investigated the possibility that reduction in the level of expression of the *FLF* gene accompanies the transition to flowering. RNA was isolated from C24 and *flf* whole plants every 10 days post-sowing, until the stage where the majority of C24 plants had bolted (50 d). The expression of the *FLF* gene remained unaltered in these plants (Figure 5C), suggesting that if reduction in the level of expression of the *FLF* gene does accompany the transition to flowering it must occur in only a few cells.

The *flf* mutant demonstrates a number of similarities to the *lhy* mutant described by Schaffer et al. (1998): they are both (semi-)dominant late-flowering mutants caused by insertion of foreign DNA adjacent to the gene. In wild-type plants the *LHY* gene is expressed for only a few hours around dawn, whereas in the *lhy* mutant, the *LHY* gene is expressed around the clock. Because of the similarities between *flf* and *lhy*, we examined the expression of the *FLF* gene in C24 and *flf* tissue harvested at dawn and dusk of an 8 h photoperiod. Although there was some difference in the expression of the genes between the

two time points, there was no alteration of this pattern in the mutant.

Example 15 The Late-Flowering Phenotype Of The *flf*
Mutant Is Suppressed By Vernalization Or By
Gibberellic Acid Treatment

In a number of late-flowering mutants and ecotypes, low temperature treatment of germinating seed (vernalization) induces early flowering (Napp-Zinn, 1985), with a 4°C treatment for 21 days saturating the vernalization requirement to produce the shortest time to flower (Bagnall, 1992). The effect of vernalization on the time to flowering of hemizygous and homozygous *flf* mutants is shown in Table 2.

Table 2

Flowering Time of Vernalised C24 and Mutant Plants

	LENGTH OF VERNALISATION		
	0 weeks	4 weeks	8 weeks
C24 (wild-type)	25.2 ± 0.2	13.6 ± 0.3	--
Hemizygous <i>flf</i>	71.4 ± 1.2	39.3 ± 3.7	20.2 ± 0.9
Homozygous <i>flf</i>	>150	100.8 ± 10.7	17.6 ± 1.3

Twenty seeds of the *flf* mutant and wild-type C24 were grown aseptically on MS medium in test-tubes, and exposed to either 4 or 8 weeks at 4°C. Non-vernalized plants were grown in soil (20 plants per 20cm pot). All plants were then grown at 23°C under fluorescent lights (16 h light, 8 h dark). The data are presented as the average number of days (\pm standard errors) until stem elongation, excluding the period of vernalization.

A 28 day treatment at 4°C resulted in a substantial reduction in the flowering time. However, eight weeks at 4°C was required to saturate the

vernalization response in both hemizygotes and homozygotes to give a flowering time similar to that of the C24 control. This implies that there is an interaction between *FLF* gene expression and a component of the vernalization-induced pathway.

RNA was extracted from 12 d old C24 seedlings that were either vernalized or were untreated controls, and probed with *FLF* gene-specific probe. Figure 5E shows a dramatic decrease in *FLF* expression in vernalized seedlings compared with unvernallized seedlings, suggesting that a component of the vernalization signalling pathway controls *FLF* gene expression. Day 1 is the day on which seeds were transferred to the growth room. In *flf* mutant plants the level of transcript was reduced in 3 week vernalized seedlings, but not to the low levels observed in C24, consistent with its only partial earlier-flowering character.

FLF mutant plants vernalized for 8 weeks had a greater reduction of *FLF* transcript, consistent with the greater reduction in flowering time.

As with other late-flowering vernalization-responsive mutants and ecotypes of *Arabidopsis*, *flf* mutant plants responded to applications of gibberellic acid (GA_3) by flowering earlier. Four week-old *flf* homozygotes treated with 1 μg of GA_3 every second day for a total of two weeks flowered two weeks after the final GA_3 application, compared to later than 20 weeks for untreated plants. However, a single treatment of 4 week old plants with 1 μg of GA_3 was not sufficient to induce flowering of *flf*, although this amount of GA_3 induced early flowering of the late-flowering *fca* mutant (Bagnall, 1992), suggesting that *flf* requires a greater amount of GA_3 for floral induction. In contrast to the dramatic effect of vernalization on the expression of the *FLF* gene, exposure of either C24 or *flf* seedlings to 10^{-5} M GA_3 had no effect on the expression of the *FLF* gene.

Example 16 Movement Of An Ac Element Present Within The
T-DNA Causes Alteration In Flowering Time

Two early flowering plants (M1 plants designated efSL3 and efSL4) were identified from one seedlot comprising bulked seed from *flf* mutant plants. Both plants flowered after 18 days, earlier than C24 which flowered after 30 days. PCR analysis using primers from within the T-DNA sequence and flanking genomic sequence confirmed that these early-flowering plants were derived from the *flf* mutant, and were not contaminants.

The tandem T-DNAs present in the *flf* mutant each contain an Ac element, and we considered the possibility that movement of Ac was the cause of the early-flowering phenotype. DNA was isolated from individual early-flowering M2 progeny of efSL3 and efSL4, digested with EcoR1 and probed with the 3' region of Ac. Figure 6A shows the appearance of a new 2.1 kb band in the early-flowering plants, indicating the movement of Ac. The maintenance of the two original Ac bands present within the T-DNAs indicates that the Ac elements have remained in their original positions as well.

In order to determine the new location of Ac, we probed a similar DNA gel blot with probe 4 (Figure 2C), revealing a change in size of the 2.7 kb EcoRI fragment containing the promoter, MADS domain and part of intron I of the *FLF* gene (Figure 6B). The size of the 3' Ac fragment (2.1 kb) indicated that the Ac element had inserted near an EcoRI site, with the 3' end of the Ac nearest the EcoRI site. PCR using a primer in the 3' end of the Ac and primers near the EcoRI sites at either end of the 2.7 kb EcoRI fragment revealed that Ac had inserted within intron I in the *FLF* gene of efSL3. Sequencing of the PCR product determined the precise insertion point of the Ac element (Figure 6C). RNA was isolated from rosette leaves of early-flowering M2 plants and comparable-sized rosette leaves of C24 and *flf* plants, and probed with an *FLF* gene specific riboprobe. As shown in Figure 5d, *FLF*

gene expression was reduced to approximately 5% of the C24 expression level. It appears that the presence of Ac reduces either the transcriptional efficiency, RNA stability or the splicing efficiency of the transcript, hence reducing the amount of the normally-spliced mRNA, and resulting in early-flowering.

Example 17 Excision of Ac From Intron I Causes Later Flowering and Increased *FLF* Transcript Levels

Twenty progeny of the M1 efSL3 plant were grown and their flowering time recorded. 15 M2 plants flowered at 18 days, the same as their M1 parent; however, 5 plants flowered later than their parent. These later-flowering plants were termed "neo-lates" to distinguish them from the original late-flowering *flf* mutant plants. Their approximate days to bolting were: 50 d (3.2), 50 d (3.4), 60 d (3.5), 85 d (3.5), 100 d (3.1). In the M2 progeny of efSL4, 36 plants flowered at the same time as their M1 parent but one neo-late plant, 4.6, flowered at 38 d. DNA was isolated from individual plants and probed with 3' Ac (Figure 7A). All the neo-lates appeared to be hemizygous for the presence of the 2.1 kb band, indicating hemizygosity for the presence of Ac in intron I. In some cases (3.1, 3.2, 3.3) the Ac element had relocated to a new site, as indicated by the appearance of the new band in Figure 7A.

To demonstrate that in each neo-late plant one copy of the Ac in intron I had excised, PCR was performed using primers derived from intron I sequence, flanking the site of Ac insertion. In each case a PCR product was generated, indicating that at least one copy of Ac had excised. The presence of the 2.1 kb band in Figure 7A indicates that at least one copy of Ac remains in intron I; thus each plant is hemizygous for the presence of Ac in intron I

The neo-late plant 4.6 is homozygous for the loss of the Ac element within the T-DNA closest to *FLF* (gene B).

- 37 -

We believe that the sector of the M1 early-flowering parent, from which the 4.6 seed derived must have been hemizygous for the presence of this Ac element, and plant 4.6 is a homozygous segregant for the loss of this Ac element.

- 5 Presumably the loss of this Ac element was an event independent of that which resulted in the insertion of Ac in intron 1, as M2 progeny of the M1 plant efSL3 are homozygous for the presence of Ac at this location.

10 Total RNA was extracted from approximately 80 d old rosette leaves of the neo-lates and the gene B expression level was compared to that of 80 d rosette and cauline leaves of *flf* and 25 d rosette leaves of C24. In all neo-lates the transcript level was higher than that of the early-flowering parent, although lower than that of the
15 *flf* mutant, consistent with them being hemizygous for the late allele.

Thus insertion of an Ac element into intron I of *FLF* (gene B) greatly reduces *FLF* (gene B) transcript levels and causes early flowering, and excision of Ac from intron
20 I restores expression of the *FLF* (gene B) transcript and results in later flowering. This provides compelling evidence that over-expression of *FLF* (gene B) in the late-flowering *flf* mutant is the cause of the late-flowering phenotype.

25

Example 18 The Expression of the *FLF* Gene is Controlled
by Known Flowering Time Genes

In order to further understand the role of the *FLF* gene in the control of flowering we examined its
30 expression in a range of *Arabidopsis* ecotypes and late-flowering mutants. Figure 8A shows the expression of the *FLF* gene in a variety of ecotypes. Interestingly the *FLF* gene is highly expressed only in ecotypes which have a late allele at the *FRI* locus (Pitztal and C24), and is
35 particularly highly expressed in C24, but not in ecotypes with an early allele at the *FRI* locus (Columbia, Ws, Landsberg erecta).

It should be noted that the Pitztal seed source used in this experiment was not the late-flowering Pitztal variety. Subsequent analysis of *FLF* expression in the late-flowering Pitztal variety revealed a very high expression level, approximately three times that of C24. This correlates well with the observation that Pitztal takes three times as long to flower as C24 (approximately 90 days compared to approximately 30 days).

To investigate this further, we looked at the expression of the *FLF* gene in the Landsberg *erecta* ecotype with late alleles of either *FRI* or *FLC* (Figure 8B). Again *FLF* is expressed in the line of Landsberg *erecta* with the late *FRI*^{Sf2} allele; however, it is also expressed in lines which contain late *FLC*^{Sf2} and *FLC*^{col} alleles. The expression of the *FLF* gene in Landsberg *erecta*-*FLC*^{col} is interesting, as this plant has the same *FRI* and *FLC* genotype (*FRI*^{early}, *FLC*^{late}) as the ecotype Columbia, yet there is expression in Landsberg *erecta*-*FLC*^{col}, but not in Columbia. This suggests that Landsberg *erecta* and Columbia differ in a third, unknown, locus, and that this locus in conjunction with late alleles at the *FLC* locus is able to induce expression of the *FLF* gene, in the absence of a late allele of *FRI*.

Many of the late-flowering mutants are in the Landsberg *erecta* ecotype, and we looked to see whether the *FLF* gene is upregulated in these mutants. Figure 8C shows that the *FLF* gene is upregulated in the *fca* and *fve* mutants and slightly upregulated in the *fpa* and *fd* mutants in the Landsberg *erecta* ecotype and in *ld* in Ws ecotype, but not in any of the other late-flowering mutants tested. *FLF* is also upregulated in the *fld* mutant in Columbia ecotype. These data demonstrate that the function of the wild-type alleles of the *FCA*, *FVE*, *FPA*, *LD*, *FD* and *FLD* genes is to down-regulate the *FLF* gene.

We have shown that the higher *FLF* transcript levels in at least one of these mutants, *fca*, is the cause of the late-flowering phenotype, by using the *FLF* antisense construct from Example 13 to decrease both *FLF* transcript

- 39 -

level in the *fca* mutant and the flowering time of the mutant. We have also shown that a 28 day vernalization period is sufficient to decrease *FLF* transcript in all 6 mutants, and to decrease flowering time. Thus cold-treatment is able to overcome the up-regulation of the *FLC* gene caused by mutations of these loci, and to overcome the delay in flowering.

We also looked at the expression of the *FLF* gene in the mutants *vrn1* and *vrn2*, which have reduced response to vernalization. Both mutants were isolated in the *fca* mutant background, and as *fca* and *vrn2* are closely linked only the *fcavrn2* double mutant is currently available. No increase in expression of the *FLF* gene in *vrn1* was detectable, however, the *vrn2fca* double mutant had increased in *FLF* gene expression over the *fca* mutant level, suggesting that irrespective of any role in the vernalization response, the wild-type *VRN2* gene acts to repress *FLC* expression and thereby promote flowering. *fca vrn1* has a similar flowering time and a similar level of *FLC* transcript to *fca*. Both *fca vrn1* and *fca vrn2* have a smaller reduction in flowering time in response to vernalization than does *fca*, and this is matched by a smaller reduction in *FLC* transcript level. This indicates that the wild-type *VRN1* and *VRN2* genes are involved in mediating the vernalization-induced down-regulation of the *FLC* gene. In our growing conditions *vrn1* segregated away from the *fca* mutant, is late-flowering and shows little vernalization response, in terms of either flowering time or alteration to *FLC* transcript level. This suggests that the *VRN1* gene may be active in an *FLC*-independent pathway as well as in the *FLC*-dependent vernalization pathway.

The expression of the *FLF* gene in early-flowering plants with reduced levels of methylation (Finnegan et al, 1996) is reduced (Figure 5F), suggesting that methylation may play a role in controlling the expression of the *FLF* gene, or a gene which is a regulator of *FLF*.

Vernalisation, or some component in the vernalisation signal transduction pathway, acts either to suppress *FLF* transcription or to increase *FLF* mRNA degradation. C24 has a strong vernalisation response, with plants vernalised for 3 weeks flowering in about half the time of unvernalsed plants. A four week vernalisation period decreases *flf* flowering time somewhat, but an eight week period is required to bring flowering back to C24 times. This incomplete effect of short vernalisation periods on flowering time of the *flf* mutant correlates with the incomplete decrease in transcript levels in *flf*. This suggests that the higher level of the *FLF* transcript, and presumably of the *FLF* protein, titrates out the promoter of flowering produced in response to vernalisation. Longer periods of vernalisation may produce more of the promoter, which might overcome the higher *FLF* transcript level.

Other ecotypes such as *Ws*, *Landsberg erecta* and *Columbia* show little response to vernalisation; under similar conditions to those which give good response with C24, the other ecotypes flower only 1 or 2 days early. We note that these ecotypes have very low *FLF* transcript levels, which cannot be decreased much further by vernalisation.

Pitztal is a late-flowering ecotype which has a strong vernalisation response, and has high *FLF* transcript levels. Vernalisation is expected to decrease the level of *FLF* transcript in these plants. Vernalization decreases the level of *FLF* transcript in this ecotype, with longer periods of vernalization resulting in a proportionally greater decrease in *FLF* transcript, correlating with the decrease in flowering time.

The *flf* mutant requires prolonged GA treatment to cause it to flower early. This suggests that the high levels of *FLF* transcript (and presumably *FLF* protein) may act to remove GA, or to decrease GA action. Since *FLF* is a MADS box transcription factor, it may do so by activating genes involved in the catabolism of GA, either directly or

- 41 -

indirectly, or by altering the expression of genes involved in GA signal transduction.

The lack of effect of GA treatment on the level of *FLF* transcript suggests that GA acts either downstream
5 of *FLF*, or via another pathway. ie.

VERNALISATION → → decrease in *FLF* transcript → → GA → → FLOWERING
or

VERNALISATION → → decrease in *FLF* transcript → → FLOWERING
10 ? → → GA → → FLOWERING

with *FLF* normally acting to block the pathway between VERN and GA

15 *Tissue-Specific Expression*

Levels of expression of *FLF* in different tissues of the C24 ecotype were examined. High expression was observed in vegetative leaves, roots, flower buds and mature flowers. There was a low level of expression in
20 cauline leaves and bolt stem, and a very low expression level in green siliques. RNA was isolated from C24 rosette leaves and "apex", ie. the tissue remaining after as many as leaves as possible and the roots were removed; this tissue includes very small leaves and the apical meristem.
25 There was no difference in expression level between these two tissue types.

In the *flf* mutant expression of *FLF* was twice that of C24 in vegetative tissue, and expression in floral tissues was relatively greater (about 3 times C24 level).

30 In the Columbia ecotype the level expression vegetative leaves was very low compared to C24, while in floral tissue there was about the same level of expression as C24.

This suggests that there may be a separate
35 control of vegetative and floral transcription of the *FLF* gene.

Developmental Expression

As *FLF* appears to be a repressor of flowering, one prediction about its pattern of expression is that it might decrease prior to, or accompanying, the transition to flowering. RNA was extracted from whole plants that were grown on MS medium and harvested every 10 days. Under these conditions 50% of C24 plants were bolting after 50 d. There was no change in the level of *FLF* transcript in either C24 or the *flf* mutant. This suggests that if there is a decrease in the level of *FLF* transcript accompanying the transition to flowering it must occur in very few cells. This result also suggest that there is no decrease in the level of transcript in older leaves, ie. the transcript is not diluted out as the leaf grows.

Circadian Response

C24 and *flf* plants were grown in 8 h fluorescent photoperiod for 21 d and either maintained in this condition, or transferred to either continuous light or continuous dark. RNA was extracted from plants harvested at either the start or the end of what would have been the 8 h photoperiod. In each case there was slightly higher expression at the earlier time point, suggesting a subtle circadian response. There was no difference in the pattern of expression between the mutant and C24.

Example 19 Isolation of a *Brassica napus* *FLF* homologue

Low stringency screening of a *Brassica napus* genomic library with an *FLF* probe lacking the MADS box region resulted in the isolation of 18 strongly-hybridizing plaques out of a total of 72,000 screened. The low stringency conditions were: hybridization at 28°C in 50% formamide, 3 x SSC, 0.1% SDS, 20 x Denhardt's, 50 µg/ml salmon sperm DNA overnight and washed with a final wash of 0.1 x SSC, 0.1% SDS at room temperature. These plaques were purified, and a selection were sequenced as described in Example 4.

The partial genomic sequence of the *Brassica napus* *FLF*-like gene is set out in SEQ ID NO. 4, and the amino acid sequence of the predicted translation product is set out in SEQ ID NO. 5. The partial genomic sequence, showing the location of exons and the sequence of the corresponding translated product, is illustrated in Figure 9, and the sequence of the predicted translated product from the *Brassica napus* gene is compared with the corresponding product from *Arabidopsis thaliana* *FLF* in Figure 10. There is a high degree of conservation, with 79% identity and 83% similarity in the deduced *FLF* protein sequence (as determined by the University of Wisconsin Genetics Computer Group software package version 9.1, using default parameters).

A cDNA library was prepared from *Brassica napus*, and 10 cDNA clones were isolated using hybridisation to an *FLF* cDNA from *Arabidopsis*. The starting RNA was isolated from leaves of *Brassica napus*, cultivar Columbus. Poly(A)+ mRNA was isolated using a mRNA purification kit (Amersham Pharmacia Biotech). A cDNA library was constructed using a SuperScript Choice System cDNA synthesis kit (Gibco BRL) in the lambda-ziplox vector (Gibco BRL) with *EcoRI* arms. The primary titre of the library was approximately 500,000 pfu. Approximately 200,000 plaques from the primary library were screened, using the *Arabidopsis* *FLF* cDNA without the first two exons of the coding region. Screening was carried out at high stringency. Thirteen plaques were picked in the first round, and of these 10 were confirmed as positive by a second round of screening at high stringency. These plaques were purified, and a plasmid containing the cDNA was excised from each clone. The complete nucleotide sequence of each clone was determined, and from this the amino acid sequences encoded by each clone were deduced. Multiple sequence alignments were used to determine the relationships between the clones. From both amino acid and nucleotide sequence data it was concluded that the clones represent transcripts from 5 different genes.

The cDNA sequences and predicted amino acid sequences are set out in SEQ ID NOS: 6 to 15 and SEQ ID NOS: 16 to 23 respectively. The sequences probably represent 5 genes, grouped as follows:

- 5 12.1/16.1
- 15.1/16.2/18.2
- 11.2
- 14.1/18.1/20.1
- 11.3

- 10 The partial genomic sequence and translation in Figure 9 correspond to the cDNAs 12.1/16.1.

Example 20
protein

Expression and Immunodetection of FLF

- 15 The FLF protein could be detected on Western blots using antibodies raised to a bacterially expressed protein.

- A truncated FLF protein lacking the first 80 N-terminal amino acid residues and possessing an in-frame N-terminal histidine tag was overexpressed in *E.coli* strain BL21 [DE3] using a pET 22b+ expression vector (Novagen). 1 ml of an overnight bacterial culture was added to 100ml of LB broth containing 50 µg/ml ampicillin and the culture grown to an OD_{600nm} of 0.6 at 37°C. 100 µl of 1M IPTG was then added, and the culture was grown for a further 4 hours before harvesting for protein isolation. Histidine-tagged protein was purified using Talon metal affinity resin (Clontech). The purified protein was injected into a rabbit for antibody production (Harlow and Lane, 1988).

- 30 Twelve day old *Arabidopsis* plants were ground in liquid nitrogen and homogenised in extraction buffer (0.1M NaPO₄ pH 7.2, 1mM EDTA). Insoluble material was pelleted by centrifugation at 16,000g for 2 minutes. The supernatant was immediately boiled in a one third volume of 6x SDS sample buffer (0.5M Tris pH 6.8, 10% SDS, 0.6M DTT, 0.012% bromophenol blue) for 5 minutes. Insoluble proteins were extracted by homogenising and boiling for 5 minutes in 2x

- 45 -

SDS sample buffer (0.167M Tris pH 6.8, 3.3% SDS, 0.2M DTT, 0.004% bromophenol blue). Protein extracts (50 µg per lane) were separated on a denaturing 12% polyacrylamide gel before blotting onto a Protran nitrocellulose membrane (Schleicher & Schuell). Blots were incubated with either preimmune serum diluted 1:1000 or with FLF polyclonal antiserum diluted 1:3000. The immunoreactive protein was visualised using the ECL western blotting analysis system (Amersham) with the secondary antibody diluted 1:2000. The blots were exposed to X-ray film (Fuji RX) for 2 to 10 minutes.

The amount of *FLF* protein was dramatically decreased following 35 days of low temperature treatment in the C24 and Pitztal ecotypes and in the *flf* mutant. The decrease in the *flf* mutant was not as great as in the two ecotypes, and this finding was consistent with our results for both flowering time and *FLF* RNA transcription. There was very little *FLF* protein in either the Landsberg erecta ecotype or in the *efSL4* loss-of-function mutant, also consistent with the RNA data. These results suggest that protein expression parallels RNA expression, and that differences seen in the RNA transcript levels will be reflected in protein levels.

These results also demonstrate that immunoassay using antibody directed against *FLF* protein can be used to identify plants having either low or high level expression of this protein, and to select strains having the desired characteristics. Such immunoassays can also be used to monitor recombinant expression of the protein in bacterial or other hosts.

Example 21 Effect of transformation with *FLF* on Flowering Time in *Brassica napus*

A cassette containing the *Arabidopsis FLF* cDNA under the control of the Cauliflower Mosaic Virus (CaMV) 35S promoter was inserted into the vector pWBVec8. This vector contains the *HPT* gene, which confers resistance to

the antibiotic hygromycin, also under the control of the CaMV 35S gene. The plasmid, in the *Agrobacterium tumefaciens* strain AGL1, was used to transform *Brassica napus* cultivar BLN1239 hypocotyl explants, and plants were regenerated. Hygromycin-resistant T₀ plants were transferred to soil. Approximately four months after the transformation experiment was begun, plants were transferred to small soil pots on a misting bench, and after approximately four weeks the plants were transferred to large pots and put in a glasshouse.

Plants were regenerated from 21 independent calli. All the plants regenerated from each callus are referred to herein as a family, and each family represents 1 or more independent transformation events.

The morphology of the plants was monitored, and the results are summarised in Table 3.

Table 3

Family	Clones	Date recorded floral (and leaf no.). Floral status recorded on Days 1, 11, 25, 33 and 41.	PCR test for transgene	Antibody test for Arabidopsis FLF protein
CS Ex 1.1	/1	Small plant	-ve	-ve
	/2	Day 1	-ve	-ve
	/3	Day 1	-ve	-ve
	/4	Day 1	-ve	-ve
	/5	Day 11	-ve	-ve
CS Ex 1.2	/1	Day 25. 17 leaves	+ve	+ve
	/2	Day 33. 15 leaves	+ve	nd
	/3	Day 11	-ve	-ve
	/4	Day 33. 17 leaves	-ve (false -ve ?)	+ve
	/5	Day 25	+ve	+ve
CS Ex 1.3	/1	Day 41. 21 leaves	+ve	+ve
	/2	Day 41. 21 leaves	+ve	+ve
	/3	Day 1	-ve	-ve
CS Ex 1.4	/1	Day 1	-ve	-ve
	/2	Day 1	-ve	-ve
CS Ex 1.5	/1	Day 1	-ve	-ve
CS Ex 1.6	/1	Day 33. 15 leaves	+ve	+ve
CS Ex 1.7	/1	Day 33. 20 leaves	+ve	+ve
	/2	Day 33. 17 leaves	nd	+ve
	/3	Day 33. 19 leaves	+ve	+ve
	/4	Day 33. 16 leaves	+ve	+ve
	/5	Day 11	+ve	nd
CS Ex 1.8	/1	Day 41. 24 leaves	+ve	nd
	/2	Day 41. 22 leaves	+ve	+ve
	/3	Day 41. 24 leaves	+ve	+ve
	/4	Day 41. 23 leaves	+ve	nd
	/5	Day 41. 24 leaves	+ve	nd
CS Ex 1.9	/1	Day 25. 15 leaves	Nd	+ve
CS Ex 1.10	/1	Day 25	+ve	-ve
CS Ex 1.11	/1	Day 25	+ve	nd
	/2	Day 25	-ve	-ve
CS Ex 1.12	/1	Day 1	-ve	-ve
CS Ex 1.13	/1	In TC	In TC	In TC
CS Ex 1.14	/1	Day 25	-ve	-ve
	/2	Day 25	-ve	-ve
	/3	Day 25	-ve	-ve

CS Ex 1.15	/1	Day 33. 15 leaves	+ve	nd
	/2	Day 33. 16 leaves	+ve	+ve
	/3	Day 33. 16 leaves	+ve	nd
CS Ex 1.16	/1	Day 33. 14 leaves	+ve	+ve
CS Ex 1.17	/1	Day 1	+ve	nd
CS Ex 1.18	/1	In TC	In TC	In TC
CS Ex 1.19	/1	Day 25. 15 leaves	+ve	nd
CS Ex 1.20	/1	Discarded	Discarded	Discarded
CS Ex 1.21	In TC	In TC	In TC	In TC

In TC = plants remain in tissue culture stage
nd = not determined

5 The plants exhibited a range of flowering times, with some plants flowering in tissue culture or while on the misting bench, while others took more than 2 months longer. Polymerase chain reaction (PCR) analysis using primers specific for the CaMV35S::*FLF* transgene was
10 performed to determine which of the putative transgenic plants contained the transgene. An antibody test (Western analysis) was also carried out to determine which plants contain the *Arabidopsis* FLF protein. The antibody does not cross-react with any *Brassica* proteins under the conditions
15 used. Although the PCR and antibody tests were single experiments, there is a good correlation between the two tests.

Table 4 summarises these results.

Table 4

Flowering Time in *Brassica napus* Plants Transformed with *FLF*

Days (Taking day of potting as day 1)	Number of floral plants (those with buds or open flowers)	Number of floral PCR or Antibody positive plants	Number of floral plants negative for both PCR and Antibody test	Number of families with vegetative plants (18 families total)
Day 1	10/41	1/26	9/15	13
Day 11	13/41	2/26	11/15	12
Day 25	24/41	9/26	15/15	7
Day 33	32/41	19/26		2
Day 41	41/41	26/26		0

5 The day of potting into large pots is referred to as day 1. Floral plants are those with buds or open flowers.

10 Although no wild-type flowering time control could be included in the experiment, due to the tissue culture stage of the experiment, it is reasonable to assume that the earlier flowering plants, which tested negative for the presence of the transgene and of *Arabidopsis* *FLF* protein, flowered at around the time of the wild-type plants. Thus some of the transgenic lines flowered at
15 least 6 weeks later than the earliest flowering lines. The delay in time to flowering was also manifested by an increase in the number of leaves which the plant developed before floral buds appeared.

20 This example demonstrates that transformation with *FLF* can be used to modify the time of flowering. The person skilled in the art will be able to apply these findings to other species of plant.

Example 22 FLF-like Molecules from Arabidopsis
thaliana

Searching of the Arabidopsis genomic sequence database revealed five putative MADS-box encoding genes with a high degree of homology to the FLF protein. One of these genes, designated *FLF-LIKE1*, occurs on the chromosome 1 BAC F22K20 (AC002291), while the other four, designated *FLF-LIKE 2*, *3*, *4* and *5* respectively, occur in a cluster on the two contiguous chromosome 5 P1 clones MXK3 and MQN23 (AB019236 and AB013395).

The amino acid sequences of the FLF-LIKE proteins are shown below, with amino acids identical to corresponding amino acids of the FLF protein depicted in bold type.

FLF-LIKE1 (SEQ ID NO:26):

MGRRK**IEIKRIENKSSRQVTFSKRRNGLIDKARQLSILCESSVAVVVVSASGKLYDSSS**
GDDISK**IIDRYEIQHADELRALDLEEKIQNYLPHKELLETVQSKLEEPNVDNVSVDSL**
S**LEEQL**ETALSVSRARKAELMMEY**IESLKEKEKLLREENQVLASQMGKNTLLATDDERG**
MFPGSSSGNKIPETLPLLN.

FLF-LIKE2 (SEQ ID NO:27):

MGRKK**VEIKRIENKSSRQVTFSKRRNGLIEKARQLSILCESSIAVLVVS**SGSGKLYKSAS
GDNMSK**IIDRYEIIHHADELEALDLAEKTRNYLPLKELLEIVQSKLEESNVDNASVDTL**
S**LEEQL**ETALSVTRARKTELM**MG**EVKSLQKTENLL**REENQTLASQVGKKTFLVIEGDRG**
MSWENGSGNKVRETLP**LLK**.

FLF-LIKE3 (SEQ ID NO:28):

MGRRK**VEIKRIENKSSRQVTFSKRRKGLIEKARQLSILCESSIAVVAVS**SGSGKLYDSAS
GDNMSK**IIDRYEIIHHADELKALDLAEKIRNYLPHKELLEIVQSKLEESNVDNVSVDSL**
S**MEEQLE**TALSVIRAKTELM**MD**MKSLQER**EKLLIEENQILASQVGKKTFLVIEGDRG**
MSRENGSGNKVPETLS**LLK**.

FLF-LIKE4 (SEQ ID NO:29):

MGRRK**VEIKRIENKSSRQVTFCRRNGLMEKARQLSILCESSVALIIISATGR**LYSFSS
GDSMAKILSR**YELEQADDLKTLDLEEKTLNYLSHKELLETIQCKIEEAKSDNV**SIDCLK
S**LEEQLK**TALSVTRARKTELM**ELV**KTHQ**EKEKLLREENQSLTNQLIKMGKMKKSVEAE**
DARAMSPSSSDNKPPETL**LLK**.

FLF-LIKE5 (SEQ ID NO:30):

MGRRRVEIKRIENKSSRQVTFCKRRNGLMEKARQLSILCGSSVALFIVSSTGKLYNSSS
 GDMAKIISRFKIQQADDPETLDLEDKTQDYLSHKELLEIVQRKIEEAKGDNVSIESLI
 5 SMEEQLKSALSVIRARKTELLMELVKNLQDKEKLLKEKNKVLASEVGKLLKILETGDER
 AVMSPENSSGHSPPETLPLLK.

Alignments of the full-length deduced amino acid sequence of the FLF-LIKE proteins with the full length FLF protein sequence revealed a 65.3 % identity (86.7 % similarity) for FLF-LIKE1, 61.2 % identity (84.2 % similarity) for FLF-LIKE2, 60.7 % identity (84.2 % similarity) for FLF-LIKE3, 60.7 % identity (85.2 % similarity) for FLF-LIKE4 and 56.1 % identity (86.2 % similarity) for FLF-LIKE5. In contrast, the published Arabidopsis MADS-box proteins which are most similar to FLF show only 42.9 % identity (66.3 % similarity) in the case of AGL14 (Rounsley et al., 1995), 40.3 % identity (75.5 % similarity) for CAL (CAULIFLOWER, Kempin et al., 1995) and 38.8 % identity (74.0 % similarity) for AP1 (APETALA1, Mandel et al., 1992). (% identity and similarity determined by the Wisconsin Package Version 10.0, Genetics Computer Group (GCG), Madison, Wisc., using default parameters).

cDNA from a chromosome 1 *FLF* -like gene (*FLF-LIKE1*) was isolated using a RT PCR based method. First strand cDNA was generated from 5 µg of Col-0 total RNA. Reactions were carried out using Superscript II (GIBCO BRL) in a 20 µl volume according to the manufacturer's instructions. *FLF-LIKE1* transcript was amplified by PCR using 1µl of the first strand cDNA synthesis reaction as template with primers :

5'-ATTGAATTCGGGCATAACCCTTATCGGAGATTTG-3' and

5'-AACGGATCCGTTGATGATGGTGGCTAATTGAGCAG-3';

Eco RI and Bam HI restriction sites respectively are underlined. The amplification reaction was carried out in a final volume of 40 µl, which contained 2.5 µM of each oligonucleotide primer, 1.0 units of Amplitaq Polymerase (Perkin Elmer) and 250 µM of each of the four

deoxynucleotides. Conditions for amplification were as follows: 94°C for 2 min, 40 cycles consisting of 15 s denaturation at 94°C, annealing at 55°C for 15 s and polymerisation at 72°C for 1 min, and a final extension at 72°C for 4 min before the temperature was decreased to 25°C. PCR products were purified using QIAquick PCR purification kit (Qiagen), digested with restriction enzymes Eco RI/Bam HI, and ligated into the corresponding restriction sites of a pBIISK+ vector (Stratagene).

Positive colonies were sequenced using universal primers with the Applied Biosystems Big Dye terminator sequencing mix according to the manufacturer's instructions, and analysed using an Applied Biosystems 377 sequencing machine (Perkin Elmer). cDNA sequences obtained were compared to *Arabidopsis* genomic sequence (BAC F22K20; AC002291). The University of Wisconsin GCG software package was employed for sequence analysis.

A binary construct containing the *FLF-LIKE1* cDNA under the control of a *CaMV 35S* promoter was generated by cloning an Eco RI/Kpn I digested PCR product into an Eco RI/Kpn I pART7 vector (Gleave, 1992) containing a *CaMV 35S* promoter. The PCR product was amplified using 200 pg of the *FLF-LIKE1* cDNA clone as template with primers:

5'-ATTGAATTCGGGCATAACCCTTATCGGAGATTTG-3' and
5'-CTAGTGGTACCGTTGATGATGGTGGCTAATTGAGC-3';

Eco RI and Kpn I restriction sites respectively are underlined. The amplification reaction was carried out as described above. The cloned PCR product was sequenced to ensure that no mutations had been introduced during the amplification procedure. The *35S::FLF-LIKE1* cassette was then subcloned into pART27 using Not I (Gleave, 1992), and introduced into *Agrobacterium tumefaciens* strain GV3101 by electroporation. Transgenic plants were generated by *in planta* transformation (Bechtold et al., 1993), using the NPT II gene as a selectable marker to identify transgenic plants.

The 35S::*FLF-LIKE1* construct was transformed into *Arabidopsis thaliana* ecotypes Landsberg erecta and C24. Twenty individual T₁ lines were selected for each ecotype, and of these about half showed a late-flowering phenotype. In the Landsberg erecta background, 12 out of 20 T₁ plants bolted 3-4 weeks post-germination, consistent with the bolting time of non-transformed wild type plants. The other 8 plants did not bolt until about 7-8 weeks post-germination. Similarly for C24 transformed lines, 9 out of 20 plants bolted about 4-5 weeks post-germination, consistent with the bolting time of wild-type C24 plants, while the other 11 T₁ lines did not bolt until 8-10 weeks post-germination. These data suggest that the *FLF-LIKE1* gene is capable of delaying flowering when overexpressed in *Arabidopsis thaliana* plants, similarly to what we have shown for *FLF*. Furthermore, it seems likely that the overexpression of the other *FLF-LIKE* genes described would have a similar effect on flowering time to that found for both *FLF* and *FLF-LIKE1* genes, especially considering that *FLF-LIKE1* shows greater homology to the other *FLF-LIKE* genes than to *FLF*.

Example 23 *FLF* Modulates Gibberellic Acid (GA)
Activity in a Number of Developmental
Processes

The phenotypic effects of over-expression of the *FLF* transcript were investigated in 35S::*FLF* transgenic *Arabidopsis thaliana*. The ecotypes used and the numbers of T₁ plants examined are indicated below. Many of the phenotypic characteristics found to be modified are known to be associated with growth processes which are controlled or modulated by GA. These include pollen formation, leaf expansion, decreased petiole angle, and trichome formation. It was particularly noteworthy that some of the plants were sterile, while others showed reduced bolt height and internode length, characteristic of a dwarf phenotype, or colour changes associated with the dwarf phenotype.

The late-flowering *flf* mutant requires much more GA to induce early flowering than wild type plants, suggesting that the *FLF* gene product may act to remove GA or GA activity. As many of the phenotypic abnormalities observed in transgenic plants expressing high levels of the *FLF* transcript can be attributed to an alteration in GA level or activity, it seems likely that this function of *FLF* is not limited to the control of GA activity in relation to the promotion of flowering, but also in relation to other roles of GA. Hence, the *FLF* gene may be useful in regulating GA activity in other aspects of plant growth, including, but not limited to, control of plant architecture and/or fertility.

As well as the effects on flowering time that have been discussed in previous examples, over-expression of the *FLF* coding sequence also produced a number of vegetative and floral phenotypes, which are outlined below.

C24 ecotype (24 T1 plants)

One early-flowering plant (#6) had the appearance of a semi-dwarf (bolt height reduced, internode length reduced), typical of the GA semi-dwarfs, caused by a reduction but not abolition of GA production in the plant. This plant was sterile.

Several plants (#6, 9, 10, 16, 19, 22, 23) exhibited partial or complete sterility. In several of these plants it appeared that the anthers had not dehisced, and no pollen was visible.

Landsberg erecta ecotype (24 T1 plants)

Two plants (#59, 63) were sterile or produced very few seeds. These plants had clear floral abnormalities, having petals of reduced size that were greenish in colour, and abnormally-shaped carpels with trichomes ("hairs"). Normally *Arabidopsis* carpels do not have trichomes.

- 55 -

The leaves of some plants which still have not flowered were reduced in size, resulting in a smaller diameter leaf rosette than normal (#40, 42, 62). Many of the late-flowering plants (#36, 37, 38, 40, 42, 62) were a darker green colour than normal. GA-deficient dwarfs are dark green. Many of the late-flowering plants exhibited regions of bulging in the leaves, suggesting that leaf expansion is not occurring equally in each direction. Many plants had a reduced petiole angle, ie. the leaves were flatter.

Landsberg erecta.-FLC^{Sf2} (17 T1 plants)

Two plants had the appearance of a semi-dwarf (#26, 28). The plants had reduced fertility. One plant (#25) produced very few seeds. This plant had clear floral abnormalities, with petals of reduced size that were greenish in colour, and abnormally-shaped carpels with trichomes.

One plant (#32) which has still not flowered has extremely small rosettes (~15 mm in diameter). Others are small, but not so extremely so (#100, 103). One of the late-flowering plants (#102) was a darker green colour than normal. Many of the late-flowering plants exhibited regions of bulging in the leaves, again suggesting that leaf expansion is not occurring equally in each direction. Many plants had a reduced petiole angle.

Landsberg erecta-FRI^{Sf2} (35 T1 plants)

One plant (#47) had the appearance of a semi-dwarf. This plant had reduced fertility. One plant (#89) produced very few seeds. This plant had clear floral abnormalities: petals of reduced size that were greenish in colour, abnormally-shaped carpels with trichomes. Some of the late-flowering plants (#48, 56) were a darker green colour than normal. Two plants (#97, 98) which have still not flowered, have extremely small rosettes (~15 mm in

diameter). Others are small, but not so extremely so (#51, 53, 78, 93, 95, 96).

Many of the late-flowering plants exhibited regions of bulging in the leaves, suggesting that leaf expansion is not occurring equally in each direction. Many plants had a reduced petiole angle.

Further phenotypic abnormalities in Ac generated early-flowering mutants

The early-flowering mutants generated by insertion of Ac into intron 1 have a reduced number of trichomes on their leaves.

It will be apparent to the person skilled in the art that while the invention has been described in some detail for the purposes of clarity and understanding, various modifications and alterations to the embodiments and methods described herein may be made without departing from the scope of the inventive concept disclosed in this specification.

References cited herein are listed on the following pages, and are incorporated herein by this reference.

REFERENCES

- An *et al*
EMBO J., 1985 4 277-284
- 5 Armstrong, C.L., Peterson, W.L., Buchholz, W.G.,
Bowen, B.A. and Sulc, S.L.
Plant Cell Reports, 1990 9 335-339
- 10 Bagnall, D.J.
Aust. J. Plant Physiol., 1992 19 401-409.
- Bechthold, N., Ellis, J. and Pelletier, G.
C.R. Acad. Sci. III, 1993 316 1194-1199.
- 15 Bevan, M.
Nucl. Acids Res., 1984 12 (22) 8711-8721.
- Brumbaugh, J.A., Meddendorf, L.R. and Grone, J.L.
20 Proc. Natl. Acad. Sci. USA, 1988 85 5610-5614.
- Chang, C., Bowman, J.L., De John, A.W., Lander, E.S., and
Meyerowitz, E.M.
Proc. Natl. Acad. Sci. USA, 1988 85 6856-6860.
- 25 Chou, M.-L. and Yang, C.-H.
Plant J., 1998 15 231-242.
- Christou, P., McCabe, D.E. and Swain, W.F.
30 Plant Physiol., 1988 87 671-674
- Crossway *et al*
Mol. Gen. Genet., 1986 202 179-185
- 35 Dean, C., Sjodin, C., Page, T., Jones, J. and Lister, C.
Plant J., 1993 2 69-81.

Devereux, J., Haeberli, P. and Smithies, O.
Nucl. Acids Res., 1984 12 387-395.

5 Dolferus, R., Jacobs, M., Peacock, W.J. and Dennis, E.S.
Plant Physiol., 1994 105 1075-1087.

Elledge, S.J., Mulligan, J.T., Ramer, S.W., Spottswood, M.
and Davis, R.W.
Proc. Natl. Acad. Sci. USA, 1991 88 1731-1735.

10 Feinberg, A.P. and Vogelstein, B.
Anal. Biochem., 1983 13 6-13.

Finnegan, E.J., Lawrence, G.J., Dennis, E.S. and
15 Ellis, J.G.
Plant Mol. Biol., 1993 22 (4) 625-633.

Finnegan, E.J., Peacock, W.J. and Dennis, E.S.
Proc. Natl. Acad. Scie., 1996 93 8449-8454

20 Fromm et al
Proc. Natl. Acad. Sci., 1985 82 5824-5828

Gleave, A.P.
25 Plant Mol. Biol., 1992 20 1203-1207

Harlow, E., and Lane, D. (1988). Antibodies: A Laboratory
Manual. (Cold Spring Harbor, NY: Cold Spring Harbor
Laboratory Press).

30 Herrera-Estrella et al
Nature, 1983a 303 209-213

Herrera-Estrella et al
35 EMBO J., 1983b 2 987-995

Herrera-Estrella *et al*

In: Plant Genetic Engineering, Cambridge University Press,
NY, 1985 63-93

- 5 Kempin, S.A., Savidge, B. and Yanofsky, M.
Science, 1995 267 522-525

Koncz, C. and Schell, J.
Mol. Gen. Genet., 1986 204 383-396

10

Koornneef, M., Hanhart, C.J. and Van der Veen, J.H.
Mol. Gen. Genet., 1991 229 57-66.

15

Koornneef, M., Blankestijn-de Vries, H., Hanhart, C.,
Soppe, W. and Peeters, T.
Plant J., 1994 6 (6) 911-919.

20

Krens, F.A., Molendijk, L., Wullems, G.J. and
Schilperoort, R.A.
Nature, 1982 296 72-74

Langridge, J.
Aust. J. Biol. Sci., 1957 10 243-252.

25

Lawrence, G., Finnegan, E.J. and Ellis, J.
Plant J., 1993 4 659-669

Lazo, G.R., Stein, P.A. and Ludwig, R.A.
Bio/technology, 1991 9 963-967.

30

Lee, I., Michaels, S.D., Masshardt, A.S. and Amasino, R.M.
Plant J., 1994b 6 (6) 903-909.

Lister, C. and Dean, C.

35

Plant J., 1993 4 (4) 745-750.

- Longemann, J., Schell, J. and Willmitzer, L.
Anal. Biochem., 1987 163 16-20.
- 5 Mandel, M.A., Gustafson-Brown, C., Savidge, B. and
Yanofsky, M.F.
Nature, 1992 360 273-277
- 10 McNellis, T. W., Mudgett, M. B., Li, K., Aoyama, T.,
Horvath, D., Chua, N.-H. and Staskawicz
Plant J., 1998 14 247-257.
- Napp-Zinn, K.
In CRC Handbook of Flowering Vol. 1. A.H. Halevy ed. (Boca
Raton, FL. CRC Press). 1985 pp 492-503.
- 15 Newman, T. et al
Plant Physiol, 1994 106 1241-1255
- Paszkowski et al
20 EMBO J., 1984 3 2717-2722
- Pietrzak, M., Shillito, R.D., Hohn, T. and Potrykus, I.
Nucl. Acids Res., 1986 14 (14) 5857-5868
- 25 Rounsley, S.D., Ditta, G.S. and Yanofsky, M.F.
Plant Cell, 1995 7 1259-1269
- Sanford, J.C., Smith, F.D. and Russel, J.A.
30 Methods Enzymol., 1993 217 483-509
- Sanger, F., Nicklin, S. and Coulson, A.R.
Proc. Natl. Acad. Sci. USA, 1977 72 5463-5467.
- 35 Schaffer, R.I., Ramsay, N., Samach, A., Corden, S.,
Putterill, J., Carré, I. and Coupland, G.
Cell, 1998 93 1219-1229

Shannon, S. and Meeks-Wagner, D.R.
Plant Cell, 1991 3 877-892.

Southern, E.M.

5 J. Mol. Biol., 1975 98 503-517.

Sung, Z.R., Belachew, A., Shunong, B. and
Bertrandgarcia, R.

Science, 1992 258 1645-1647.

10

Valvelkens, D., Van Montagu, M. and Van Lijesbettens, M.
Proc. Natl. Acad. Sci. USA, 1988 87 5536-5540.

Weigel *et al*

15 Cell, 1992 69 843-859

CLAIMS

1. An isolated nucleic acid molecule comprising a MADS box, which is capable of altering the flowering time of a plant.
- 5 2. An isolated nucleic acid molecule according to Claim 1, in which expression of the nucleic acid molecule in the plant, in the sense orientation under the control of a promoter sequence, is capable of delaying the flowering of the plant.
- 10 3. An isolated nucleic acid molecule according to Claim 1, which is capable of accelerating the flowering of a plant.
4. An isolated nucleic acid molecule according to Claim 3, in which expression of the nucleic acid molecule in the
15 plant in the anti-sense orientation under the control of a promoter sequence is capable of accelerating the flowering of the plant.
5. An isolated nucleic acid molecule according to Claim 1, which comprises a nucleotide sequence corresponding to
20 a *FLOWERING LOCUS F* (*FLF*) gene, or a PCR primer or a biologically active fragment derived therefrom.
6. A nucleic acid molecule according to Claim 1, which comprises
 - 25 (a) the nucleotide sequence set out in any one of SEQ ID NOS. 1, 2, 4, and 6 to 15;
 - (b) a nucleic acid molecule capable of hybridizing thereto under at least low stringency hybridization conditions; or
 - 30 (c) a nucleic acid molecule which has at least 70% sequence identity with a sequence set out in (a).
7. A nucleic acid molecule according to Claim 6, in which the nucleic acid molecule is
 - 35 (a) capable of hybridizing to a nucleotide sequence as set out in any one of SEQ ID NOS: 1, 2, 4, and 6 to 15 under high stringency hybridization conditions; or

(b) has at least 80% sequence identity with a sequence set out in Claim 6(a).

8. A vector comprising a nucleic acid molecule according to any one of Claims 1 to 7.
- 5 9. A plant cell transformed with a nucleic acid according to any one of Claims 1 to 7.
10. A plant transformed with a nucleic acid molecule according to any one of Claims 1 to 7.
- 10 11. A method of isolating a nucleic acid molecule capable of altering the flowering time of a target plant, comprising the step of using a nucleic acid molecule according to any one of Claims 1 to 7, or a functional portion thereof, as a hybridisation probe or polymerase chain reaction (PCR) primer, and optionally detecting
15 hybridisation.
12. A method according to Claim 11, in which the nucleic acid molecule is capable of hybridizing to a nucleotide sequence as set out in any one of SEQ ID NOS: 1, 2, 4, and 6 to 15 under at least low stringency hybridization
20 conditions.
13. A method of delaying flowering in a plant, comprising the step of introducing a nucleic acid molecule according to any one of Claims 1 to 7 into cells of the plant, optionally such that expression of the nucleic
25 acid molecule is under the control of an inducible promoter, and over-expressing the nucleic acid molecule.
14. A method of inducing early flowering in a plant, comprising the step of reducing the degree of
30 expression of a nucleic acid molecule according to any one of Claims 1 to 7 in the plant.
15. A method of modifying the vegetative and/or floral phenotype of a plant, comprising the step of increasing the level of expression of an *FLF* gene, thereby to
35 modify the level of production or activity of a gibberellin in the plant.

16. A method of modifying the response of a plant to vernalisation, comprising the step of increasing or decreasing the level of expression of an *FLF* gene.
17. A method according to Claim 15 or Claim 16, in which the *FLF* gene comprises a nucleic acid molecule according to any one of Claims 1 to 7.
18. A polypeptide encoded by a nucleic acid molecule according to any one of Claim 1 to 7.
19. An *FLF* polypeptide, comprising the amino acid sequence set out in any one of SEQ ID NOS: 3,5, and 16 to 30, or having at least 70% sequence identity thereto.
20. An antibody directed against a polypeptide according to Claim 18 or Claim 19.
21. A method of assaying the level of expression of *FLF* polypeptide, comprising the step of using an antibody according to Claim 20.
22. A method of selecting plants with low or high levels of expression of *FLF*, comprising the step of determining the level of *FLF* mRNA or *FLF* polypeptide in the plant.
23. A method according to Claim 21, in which the plants are members of a naturally-occurring population.

1/8






Figure 1

THIS PAGE BLANK (USPTO)

2/8

*Figure 2*

THIS PAGE BLANK (ISPTO)

*Figure 3*

THIS PAGE BLANK (ISPTO)

4/8




Figure 4

THIS PAGE RI ANK (11SP70)





Figure 5


THIS PAGE BLANK (USPTO)

6/8

*Figure 6*

THIS PAGE BLANK (ISPTO)

7/8

*Figure 7*

THIS PAGE BLANK (USPTO)

8/8




Figure 8

THIS PAGE BLANK (USPTO)

SEQUENCE LISTING

(1) GENERAL INFORMATION:

- 5 (i) APPLICANT:
(A) NAME: COMMONWEALTH SCIENTIFIC AND INDUSTRIAL
RESEARCH ORGANISATION
(B) STREET: LIMESTONE AVENUE
(C) CITY: CAMPBELL
10 (D) STATE: ACT
(E) COUNTRY: AUSTRALIA
(F) POSTAL CODE (ZIP): 2612

15 (ii) TITLE OF INVENTION: CONTROL OF FLOWERING

(iii) NUMBER OF SEQUENCES: 3

- (iv) COMPUTER READABLE FORM:
(A) MEDIUM TYPE: Floppy disk
20 (B) COMPUTER: IBM PC compatible
(C) OPERATING SYSTEM: PC-DOS/MS-DOS
(D) SOFTWARE: PatentIn Release #1.0, Version #1.30 (EPO)

25 (2) INFORMATION FOR SEQ ID NO: 1:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 7968 base pairs
(B) TYPE: nucleic acid
30 (C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

35 (iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

- (vi) ORIGINAL SOURCE:
40 (A) ORGANISM: *Arabidopsis thaliana*

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 1:

45 GGATCCAAGA AATAATTTCA TATGGGGCAC AGTTAAAAAA AAAACAATAA
AATGATAATA 60

GTAAGGCTTG AACTTGGGTT GATGTGAGGC ACTATTAAGT AAAAAGCCAT TGTACTACTT
120

50 ACATTTTAAC TACATAATGT TAACTTATAT AATATTTATT GAATTCAGTA TAGGGCACAT
180

GCCCTATCCA TGACTAACGT GAGTCCGCCC TGATAGCGAG TAAGAAACGA
55 GCAAAGGAAT 240

GCAAATTATG TGAATACTAT ACACCAACAG TGTAGACATG TAGCTACAAT GCGGCAATGT
300

THIS PAGE RI ANK (11CDT0)

TAAAAATAAA CTCAAATTGG TTTGGAGGGA ACAACCTAAT GCTTATAAGT ACACTTTTGT
360

5 GGTAATACA ATCCAATTGA AAGTCTTTGT AGGTTTGGTT TGGTCCAATG AAATTGTATG
420

CGAGTTGAGT AAAATAACCT TAGTTCAAAA CATTAGATAT GTAATGGTCT AGATACGATG
480

10 GTAGCCAAAG ATTTGGGTTA AATTTAGAAT AAAGATCAAA GAACAAATGA CTGACTTCCT
540

TATGTGTGTT TGTTTATGTA AAGTCTTAAC TCGTGTCTTG CCAAATTAAT AAAAAGGTGC
15 600

ATTATACAAT GAACTAGAGC TTGTTCTCAT CAAAATTTTT AACATTAAAA TAAGTTATTA
660

20 AAATTTTGTT GATTATATAT GATTTCAATA TTAAATTATA TTTTTTGCTT ATAGCATCAA
720

AACTTCTTGG CACAGCTCCG AGTGTTACTG AAATGTTTGT GTGGCTCCAA TAGAAAAGTT
780

25 AATACCAATC ATGAATCATC AATATCGTTA CAAAATCAGT AATCAGTTTC ACCCACTCCG
840

AGTGTTAGTG AAATGTTTGT GTGGCTCCAT TAAAAAAGTT AATACCAATC ACCAATATCG
30 900

TTACAAAATT TGTAATCAGT TTCACCCACT TCTTTTCGTC TTGTTACTCG ATATAGCCTC
960

35 ACCTTTCAAG TGTCTAATAT AACACTTTTT AACAAAATAT TACAAAACAA GAAAAATTAT
1020

AGTAATTAGA TTAATCATAT GTAATCTATT ACATATAGTA ATTTGTATTA GTATCGTTTA
1080

40 TTGTGTTACC ATTCAAACGG TATAATCTAT ATAATTATTA GCAAAACAAA ATATAGTTTT
1140

AGTAATTAAT AAAATTAATA TAATGATAGT GATATTCAGA ATGTGGTATC TAATCATGTA
45 1200

AAAATATATA ATAATAAAAT TAGATAAAGA AGAAATTGGT AAAAAATAAT
TAATGAGATA 1260

50 TAAAAGGAAA ACAAGCTGAT ACAAGCATTT CACCCAAAAA AAAACAAGCT
GATACAAGCA 1320

AAAAAGAATT AGTAACTTTG AGCTATTGCC ATATGTGTGG ACATTTAAGA TTTCTTGTAT
1380

55 TTTAGAAATC TTATATATTT CCACAATATA TTTACTACTT TCCTTTATTA TTTGTGTTAA
1440

THIS PAGE RI ANK (ISPTO)

TCTCCCGAAC ATTATTATTT CAATACTATC TGAAAAACAC ATTTTTTTTT ATAAAATTTG
1500

5 ATGACGTAGG CGAGTGGTTC TTTGTTTTTA CTATGTAGGC ACGACTTTGG TAACACCTAC
1560

TAATTAAGT CCAAATTTTA AGTTTTGAGA AGTCGGAAGA GTTCAAACCA GTTTTAGGTT
1620

10 TCGATATGTC TACATAGTTC AAAGATGATG TAGAGTGGAG GTTCTTTCTG CAATAGTTCA
1680

ATCCGTATCG TAGGGGAGGA AAGATAGTTT TCATTTAGCA AAGAAAGTGA
AAACTAAGGC 1740

15 AATGCAAAAG TAGCAAAGAC GCTCGTCATG CGGTACACGT GGCAATCTTG TCTTCAAAAC
1800

GCAACGTTTT TATTCACATA TTTGGTTTTT TTGCATCACT CTCGTTTACC CCCAAAAAAA
20 1860

AAAAATATCT GGCCCGACGA AGAAAAAGTA GATAGGCACA AAAAATAGAA
AGAAATAAAG 1920

25 CGAGAAAAGG AAAAAAAAAA ATAGAAAGAG AAAACGCTTA GTATCTCCGG
CGACTTGAAC 1980

CCAAACCTGA GGATCAAATT AGGGCACAAA GCCCTCTCGG AGAGAAGCCA
TGGAAGAAA 2040

30 AAAACTAGAA ATCAAGCGAA TTGAGAACAA AAGTAGCCGA CAAGTCACCT
TCTCAAACG 2100

TCGCAACGGT CTCATCGAGA AAGCTCGTCA GCTTTCTGTT CTCTGTGACG CATCCGTGCG
35 2160

TCTTCTCGTC GTCTCCGCCT CCGGCAAGCT CTACAGCTTC TCCTCCGGCG ATAAGTACGC
2220

40 CTTTTCTTA CCTGGGTTTT CATTTATTCC CCCTTTTATC TTCTGTTTTG TGCTCTTTA
2280

CTTTTCTGAG AAAATAAAAA TAAAAAACA ATTAATATAC CGTTTGGTTT TTTCCGGCG
2340

45 GATCTCTTGT TGTTTCTCGG TTCTGTGTTT GTTTGTGTTT TTTTCTGCGA CCATGATAGA
2400

TACATGAGAT AACCAAATTT AAGGAAGAAC AATGTCGTGA AGAAGCTTTT TAGCTTCTTA
50 2460

CTTTTGTTC TTTCTCTCTC TATTTCTTAA AAAAAAAAT TCTGCATGGA TTTCATTATT
2520

55 TCCTTGGAAG AAAATTGCAT GTCCTTCACG ATTTGTTTGA TACGATCTGA TCGGTGCTCG
2580

THIS PAGE RANK 1115DT01

ATGTTGTTGA GTGAAGTTTC AAGCCATCTT TGATTGTTTC TTACCTTTAG AGATTCCTTA
2640

5 AGTTTTTGAA GAGTTAATTA TATATCACAA GACTAATGAT TAATGTCTCT TTTCAGAGTG
2700

ATTAATAATTC ATTGGATCTC TCGGATTTGT ATGCAATGCA CTTACGGGAG ATCTATAGAG
2760

10 TTGCTATGGG GTTAATGCTG AACAATGTAT ATATACCACA TTGTGCAGCT ATTGACTATA
2820

TAAGACTATT GTTAATCTTC TATGAATTCC TATCTTTGCT GTGGACCTAT TACTTGGTGA
2880

15 TTATCCAAAT TAGTGTTTTT AATTGATTCA TATTTTTCAT ACACAGTAGT TTTGAATTTT
2940

GGTAGCTTCA AAAAATCAG CCTCACAATT AGTACTTACC GCACATATGC TACTTCAGTA
3000

20 ACATAATCTG GTTATCGATT GCGATTCTTT GAATCACAAT CGTCGTGTGC TATATATATA
3060

25 ACACCTTTTG CTGTACATAA ACTGGTCTAA TTTAGACTA ATTAAATTTT ATTGTTCTCT
3120

TGGATTTGTA TATGCACGTC CGGGAGATTT ATAAATAAAA TTAGTATGAG GTTAATGGTA
3180

30 AAAAGGATCA AGAAGTTTGG TTTTAAATGT AAGCCACATT AATTGGGAAA CTATGACTAA
3240

AAGATGAATT GGTAATATAT ACATAATATT TTTAACGAAT TTCTCTCCTT TTTATGGGAT
3300

35 ATGCTATTTG AAGATTGTCC AAAGGTTTAT AGTTTCCCAC TCTTGCAGTT ACACACATAG
3360

40 ATTTGCCTCA TATTTATGTG ATTGTATATC AATTATCGCC CTTAATCTTA TCATCGTTGT
3420

GTTCAATTAT GACTTTGTTC CTATTCGTTA AAATTGACAA TCCACAACCT CAATCTTTTG
3480

45 TTGTGAAAAT CGACAATCAC ACAACCTTTG TATCTTGTGT CTTTTGTCAT GGAAATTGTC
3540

ATTCACACAG CCTTGTTTCT TTGGTGCCTC TAGGAAATTG AAAATCCCAC AACACTTGTC
3600

50 TTCATGTAAG AAATACCAAC CTCTTTGGTA CGGATCTATA ATGAATCAAT ATAATCCTAT
3660

55 ATATAAGTTG TCAAAATTGA ATCTGGTGTA GTGTCTACTA CAACCCTCCA ATATAATAAC
3720

THIS PAGE RI ANK 1110000

CAAATGGTTG TAGTAGTTTG GCCATGTTGG TCAAGATCGC TGGCCGATTC TCACTTGATG
3780

5 CATACTTTGT TAGGATTTGT TCACCCCTAG TTAGGTCCAG CCTTGGAATT GTCGAGACAC
3840

CTGACTAGAA CTCCTGGTCT TAATTATGAT TTAATAAAGA AGAAGCCTTT TAGAACGTGG
3900

10 AACCCTTAGT TACTCAGTTA CTCTTTTTGC ATACTTAGGT TGATGCAAAG AGCTTAACTT
3960

CACAATAGGA CTGATATCTA TTAACAAAAC AAATTAAGTG AAGTTTTGTC AAAATTGTTG
4020

15 GATCTTCTAG GTCAATATGT AGTTTAGTTT TTATCTGTCT TAGTCGCTTC CTTCTATGGA
4080

AGATATATTT ATAGATATGT GTGATAAGTT TCCTACTAAT ATTAGTTAGT GTAACCTCAA
4140

20 GGGCAGAAAA CTCTTTACTT TTATTGCTTG ATTAATTTGG GGTTTAAATA TAGGAAATTG
4200

25 GAACCTCACA GTTCTTATAA ACGAGTATGT AATTGAATGT GAAATAACAA AATGGAAGAC
4260

CGGCTTCCTA TTCTTAGGAG TCTTTTGATA TTTGCAAAAA AACATATAC AATAGAAATA
4320

30 TGAGTTTTGT CTAAGACTCG GTCCATGTAT TTGGAGTTTG GCTTCCTCAT ACTTATGGTT
4380

ATCTGGTTAC CGCCACATCA TCATTATCAT CTTATGGGTC ATCAATACTA GCTCTATCGC
4440

35 TGGAAAAAAC CTTGTCCTCA AGGTTTATTG AAAAATCCGA AAAGTTTTCT CGTATATGTT
4500

40 GATATGGTAT TACTTACAAA CAAAGAGCTG ATGTTACCAA TTTTGACACG AGATTACTAA
4560

TGAACATG AAAGAGGCGT TTTTAAAAAA TTCTTTTAA AACTGGGATA CAAAAAGAAA
4620

45 AGAGGTAAC AATAATTGA TACCATTGTT CGTAGTCCTG ATCAAATGTT ATAAGGGTAA
4680

ACATGATAGA AAATAGAGGG TAAATAGGTT TTGTTCTTAT AATGGTTTTG ATAACACGCT
4740

50 TTGTAAAGGA TATAGGTGTT TTTTGATGCT AAAAGTTGTG GTATGGATCA AAACCAAAT
4800

55 GGAAGCTCTG AATCTCTGAT AGAGGTTGCA ATTAGAATTA TATAAGTTAA TTTGCAAATG
4860

THIS PAGE BLANK (15PTO)

AATTGGAAGC AGTCTTCCAC TATTTGCTAT TGTTAGGGAA GTCTTTCAGT TAATTTTCAGA
4920

5 AAATTAAGAG AAATATGACT TTCTAGACTC AGTCTGTGTA CTTGGAATTT TACTTCGGTT
4980

TACTTCCATG TCATCACATT GTGGCTCATC AATATATGTG TGTATATAÇA TTCATGAGTA
5040

10 TATATGATTT CTGGAAAAAT AAAAATTGCT TGTTTGCATT TAAGATTGGG GCTGCGTTTA
5100

CATTTTATAT TGCATCAATT ATTTCAACAT AGATTCACAA ACATAAATGC ATAGAAACAA
5160

15 TCTGGACAGT AGAGGCTTAT GTTTAGGGTT CTTATGTACC TTAAGTAGTT TGACTTTAAG
5220

TTAATCAAAG CCAGCGCTAT CACTAACTT TATCTGTATG CCTTTGTATG ACTTTTCTTT
5280

GAGGGAAAAT GTCATTTTCA ATCTGCCGAA ATATATAATA AATACATGTT AGCCACATA
5340

25 ATTCATTGGA TAACTAATCT TTGAGCAATT TTTGGTAAAT GTTTTGGTTC TTTTCTTTTC
5400

TTGAGAGAGA AAAAAAATAT CAGATATTAT TAAATATTGC TTACAAAGCT AAGAACAAGT
5460

30 TAAAACTTTT TTGAAAAAGT GGAAATTCAG ATGTGCTACT GCTTAAACAT GAATATTAAG
5520

ATTATTGTTT TTCTGAAATG TTACGAATAC TAGCGTGTTA TATATATGTA AAAGGTAAGG
5580

TGTTCTCTCA ATGTTTCATA GTTCCAGTG GCCTTTTCAA GGGTAGCTA GTAGTTTGA
5640

40 TCCTAACATA TTTTATTTT TTTTGTATC TCTCCAGCCT GGTCAAGATC CTTGATCGAT
5700

ATGGGAAACA GCATGCTGAT GATCTTAAAG CCTTGGTAAT ACAAACATTT TGAATCTTTT
5760

45 CCCTGATGGA GTTTTATAAG GCGTAAATTT ACTATTAGTT TGCCGAGTGA TCCTAAATAT
5820

AAAATGAGGT GGTGGCTCCA CATGCATTAT GCATACCGCA ATTTTCATAG CCCTTGTCTT
5880

TTACCGCTTC TTCTGTCCCT TTTTCATGGG CAGGATCATC AGTCAAAAGC TCTGAACTAT
5940

55 GGTTCACACT ATGAGCTACT TGAAGTTGTG GATAGGTTAG TACTACTAAC TAAGACTATA
6000

THIS PAGE BLANK (USPTO)

TTTGCTCTCC ACCTTTGATT ACAAAGGAAT TAGTTTTTTT TTTGTCAAAC TATGAATATA
6060

5 TGCAGCAAGC TTGTGGGATC AAATGTCAAA AATGTGAGTA TCGATGCTCT TGTTCAACTG
6120

GAGGAACACC TTGAGACTGC CCTCTCCGTG ACTAGAGCCA AGAAGGTAAG TTGATTTCGT
6180

10 AATGTCTACT CCTTTCTGAA TTTTGTGTTGC TGAGAACAAC CGTGCTGCTT TTGTTTGTG
6240

CAGACCGAAC TCATGTTGAA GCTTGTTGAG AATCTTAAAG AAAAGGTCAG ATATTTGCTA
6300

15 CCAATTTTAT TGTACATCAG ATATATCCTC TTCTGTGTTG TCTCTGTTAC TTTAAGTCTG
6360

CTTAACGAGC TTGCACACAT ATTTGCAACT TTCTTCATAT GTTTTGGATT CCAAATTCTG
6420

20 AAGTTGTTAG GTTTAGAAAC TTGATCGGTA ATTGCTGAAC ATTTTGATCT TTAAATCAGG
6480

25 AGAAAATGCT GAAAGAAGAG AACCAGGTTT TGGCTAGCCA GGTAACGAAA
GCTACATTTT 6540

CTAAAAATAT ATATGCATAA CTAATAAGCA CTGCGTGTTG TGTGTCCAAT GTCCATGTAC
6600

30 ATGGACATAG ATACACACTC TTATGCTTGC AGATATATAT ATATATATAT ATAGTCAGTG
6660

CATTTCAATC ATTCACTAGT TAGCACTTTC CTGGTCTTGT ATAGTTGTAT TCTAGACAAT
6720

35 TCTTCTCAAG ATTAGGGCAT TTTGGTTGTT GGTAGTTTGG TTTATTAGGG TTAGTGAGAT
6780

40 TATTACTGAA TAAGAACAGA AATTTGATAA CGGCTGGTTA GAGTTAAGGG AAATCAGATG
6840

AAGTTATTTT TTTATTTTTT ATCGAGTATA AATTACATGA TTGCTATATC ATTTTACTAA
6900

45 ATTAAGAAAA AAAAATTCCG GTTGTTGGAC ATAAGTAGGT TTTGGTTCTT CTCTTCGTT
6960

TTTTTCATGT TAAAGTGTTT AATTAGGTTT TGGTTCAATT GGAGATTTAG GAACCTTTTA
7020

TAGTCTGGTT AAGTCTGGGT TTGGTAGAGA TTCAATAAGA TTTCTTGATT CTCTTCAGGT
7080

55 TATGGTCTGG TTCAGTCTAG TTTAGTTCAA TATTGGTTTC CTTGAAGGTT GTGTAAACGT
7140

THIS PAGE RI ANK 1115PTO

TGTCTATATT TAAGTTAATC ACCTTTTAAC CAAAAAAAAA AGTTTATGGA CCGATTAGTT
7200

TTTTTTTTTT TTTTTTTTTG TGATGGTTAG GTTTGGATCC GAGTGGCTCA GTTCCAACCTC
5 7260

CAAGTGTCTA GAAGTAGTGC TACTTTTACA TGCTATATAT AGGTTAGATT ATAAATTATA
7320

10 AACTGGTAAA AGATTATAGA TACTGCTTCC AAACCTTAAAA GCTTAAACAT AAAGAACACA
7380

CAAATTATGA GAAAAATAAC CTTCTGTAGT GTTTTTTAAT GGTTGTTATT TGGTGGTGTG
7440

15 AAAAAGATAT TCCTTGGATA GAAGACAAAA AGAGAAAGTG AATAGTGATT
TTGACCTATG 7500

ATTATCGTAC AGATGGAGAA TAATCATCAT GTGGGAGCAG AAGCTGAGAT
20 GGAGATGTCA 7560

CCTGCTGGAC AAATCTCCGA CAATCTTCCG GTGACTCTCC CACTACTTAA TTAGCCACCT
7620

25 TAAATCGGCG GTTGAAATCA AAATCCAAAA CATATATAAT TATGAAGAAG
AAAAAAAAAT 7680

AAGATATGTA ATTATTCCGC TGATAAGGGC GAGCGTTTGT ACATCTTAAT ACTCTCTCTT
7740

30 TGGGCAAGAG ACTTTGTGTG TGATACTTAA GTAGACGGAA CTAAGTCAAT ACTATCTGTT
7800

TTAAGACAAA AGGTTGATGA ACTTTGTACC TTATTCGTGT GAGAATTGCA TCGAGATCTT
35 7860

GAGTGTATGT GTTCTTCTCT TCTGTCAAAA ACTTGTGTTT GCTTCACAGT GAAGAAGCCT
7920

40 ACGGCTTATT TTGCAACAGG GACGTGGCTC TCTCTCTCTC TCTGCGCG 7968

(2) INFORMATION FOR SEQ ID NO: 2:

- 45 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 943 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear
- 50 (ii) MOLECULE TYPE: cDNA
- (iii) HYPOTHETICAL: NO
- 55 (iv) ANTI-SENSE: NO
- (vi) ORIGINAL SOURCE:
(A) ORGANISM: *Arabidopsis thaliana*

THIS PAGE BLANK (USPTO)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 2:

5 CGAGAAAAGG AAAAAAAAAA ATAGAAAGAG AAAACGCTTA GTATCTCCGG
CGACTTGAAC 60

CCAAACCTGA GGATCAAATT AGGGCACAAA GCCCTCTCGG AGAGAAGCCA
TGGAAGAAA 120

10 AAAACTAGAA ATCAAGCGAA TTGAGAACAA AAGTAGCCGA CAAGTCACCT
TCTCCAAACG 180

TCGCAACGGT CTCATCGAGA AAGCTCGTCA GCTTTCTGTT CTCTGTGACG CATCCGTCGC
240

15 TCTTCTCGTC GTCTCCGCCT CCGGCAAGCT CTACAGCTTC TCCTCCGGCG ATAACCTGGT
300

CAAGATCCTT GATCGATATG GGAAACAGCA TGCTGATGAT CTTAAAGCCT TGGATCATCA
20 360

GTCAAAAGCT CTGAACTATG GTTCACACTA TGAGCTACTT GAACTTGTGG ATAGCAAGCT
420

25 TGTGGGATCA AATGTCAAAA ATGTGAGTAT CGATGCTCTT GTTCAACTGG AGGAACACCT
480

TGAGACTGCC CTCTCCGTGA CTAGAGCCAA GAAGACCGAA CTCATGTTGA AGCTTGTTGA
540

30 GAATCTTAAA GAAAAGGAGA AAATGCTGAA AGAAGAGAAC CAGGTTTTGG
CTAGCCAGAT 600

GGAGAATAAT CATCATGTGG GAGCAGAAGC TGAGATGGAG ATGTCACCTG
35 CTGGACAAAT 660

CTCCGACAAT CTTCCGGTGA CTCTCCCACT ACTTAATTAG CCACCTTAAA TCGGCGGTTG
720

40 AAATCAAAAT CCAAAACATA TATAATTATG AAGAAAAAAAA AAATAAGATA
TGTAATTATT 780

CCGCTGATAA GGGCGAGCGT TTGTATATCT TAATACTCTC TCTTTGGCCA AGAGACTTTG
840

45 TGTGTGATAC TTAAGTAGAC GGAACCTAAGT CAATACTATC TGTTTTAAGA CAAAAGGTTG
900

ATGAACTTTG TACCTTATTC GTGTGAGAAA AAAAAAAAAA AAA 943

50

(2) INFORMATION FOR SEQ ID NO: 3:

- 55 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 196 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: single

THIS PAGE RI ANK (11SP70)

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

5 (iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

10 (A) ORGANISM: *Arabidopsis thaliana*

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 3:

Met Gly Arg Lys Lys Leu Glu Ile Lys Arg Ile Glu Asn Lys Ser
15 1 5 10 15
Ser Arg Gln Val Thr Phe Ser Lys Arg Arg Asn Gly Leu Ile Glu
20 20 25 30
Lys Ala Arg Gln Leu Ser Val Leu Cys Asp Ala Ser Val Ala Leu
35 40 45
Leu Val Val Ser Ala Ser Gly Lys Leu Tyr Ser Phe Ser Ser Gly
25 50 55 60
Asp Asn Leu Val Lys Ile Leu Asp Arg Tyr Gly Lys Gln His Ala
65 70 75
Asp Asp Leu Lys Ala Leu Asp His Gln Ser Lys Ala Leu Asn Tyr
30 80 85 90
Gly Ser His Tyr Glu Leu Leu Glu Leu Val Asp Ser Lys Leu Val
95 100 105
Gly Ser Asn Val Lys Asn Val Ser Ile Asp Ala Leu Val Gln Leu
35 110 115 120
Glu Glu His Leu Glu Thr Ala Leu Ser Val Thr Arg Ala Lys Lys
40 125 130 135
Thr Glu Leu Met Leu Lys Leu Val Glu Asn Leu Lys Glu Lys Glu
140 145 150
Lys Met Leu Lys Glu Glu Asn Gln Val Leu Ala Ser Gln Met Glu
45 155 160 165
Asn Asn His His Val Gly Ala Glu Ala Glu Met Glu Met Ser Pro
170 175 180
Ala Gly Gln Ile Ser Asp Asn Leu Pro Val Thr Leu Pro Leu Leu
50 185 190 195

Asn

55 SEQ ID NO 4
5
6 to 25

THIS PAGE BLANK (11/15/70)

SEQUENCE LISTING

<110> CSIRO

<120> CONTROL OF FLOWERING

<130> FP10512

<140> PP7469

<141> 1998-12-03

<160> 27

<170> PatentIn Ver. 2.1

<210> 1

<211> 908

<212> DNA

<213> Brassica napus

<400> 1

```
aaaagaaaga aataaaagca aaaaagagag aaaataaaag caaaaataag aaagaacaaa 60
aaacgctcag tatctccggc gagagctgaa ccgaaccgaa cctcaggatc aaattagggc 120
acaaagggtt ctccggagaca gaagccatgg gaagaaagaa actagagatc aagcgaattg 180
agaacaaaag tagccgacaa gtcaccttct ccaaacgacg caatgggtctc atcgagaaaag 240
ctcgtcagct ttcagttctc tgcgatgcat ccgtcgtctc cctcgttgtc tcagcctctg 300
gcaagctata caacttctcc gccggcgatg acctgggtcaa gatcgttgat cgatatggaa 360
aacaacatgc tgatgatcgt aaagctctgg atcttcagtc agaagctccg aagtatgggt 420
cacaccatga gctactagag cttgtcgaaa gtaagcttgt ggaatcaa at tctgatgtaa 480
gcgtcgattc cctcgttcag ctggagaacc accttgagac tgccctctcc gtaactagag 540
ctaggaagac agaactattg ttgaagcttg ttgatagcct caaagaaaag gagaaattgc 600
tgaaagaaga gaaccagggt ttggctagcc agatggagaa gaataatctt gcgggagccg 660
aagctgataa aatggaagtg tcacctggac agatctctga catcaattgt ccggtaaactc 720
tcccactgct ttattagccc cttaagtcca aaacttgtga ctaaaaacaa aaataagtta 780
ccgaactatt cccctataag ggtgagcgtt gtatatcttc acactctctt ggctgagaga 840
ccctgtgtgt aaaaactacg gtttgatttg agtaaaaata tatatttaag aaaaaaaaaa 900
aaaaaaaaa                                     908
```

<210> 2

<211> 847

<212> DNA

<213> Brassica napus

<400> 2

```
ggcgacttta accgtacctc agaatcaa at tagggcacag agacctctcg gagacagaag 60
ctatgggaag aaaaaaacta gaaatcaagc gaatcgagaa aaacagtagc agacaagtca 120
ccttctgcaa acgacgcaac ggtctcatcg agaaagctcg tcagctttct gttctctgcg 180
```

THIS PAGE RI ANK (11CPTO)

aggcatctgt tgggcttctc gttgtctccg cctccgācaa actctacagc ttctcctccg 240
gggatagact ggagaagatc cttgatcgat atgggaaaaa acatgctgat gatctcaatg 300
ccctggatct tcagtcaaaa tctctgaact atagttcaca ccatgagcta ctagaacttg 360
tggaagcaa gcttgtggaa tcaattgatg atgtaagcgt ggattccctc gttgagctag 420
aagatcacct tgagactgcc ctctctgtaa cttagctcg gaaggcagaa ctaatgttaa 480
agcttggtga aagtctcaaa gaaaaggaga atctgctgaa agaagagaac cagggttttg 540
ctagtcagat tgagaagaaa aatcttgagg gagccgaagc tgataatata gagatgtcat 600
ctggacaaat ctccgacatc aatcttctcg taactctccc gctgcttaat taaccacctt 660
tactcggcgg ttaatcaaaa taagaaacat ataactctaaa gataacctat gtaggtttta 720
cttttcgcag cttaattaac cncctttact cggcgggttaa tcgaaattaa aaacatataa 780
ttaacaaata acctatgtca gtttaacccc aaaaaaaaaa aaaaaaaaaa gaaaaaaaaa 840
aanaaaa 847

<210> 3

<211> 969

<212> DNA

<213> Brassica napus

<400> 3

ctctggatca aattagggca cagagaccac ttggagacag aaaccatggg aagaaaaaaaa 60
ctagaaatca agcgaattga gaacaaaagt agccgacaag tcaccttctc caaacgacgc 120
agcgttctca ttgagaaagc tcgtcagctt tctgttctct gcgatgcac cgctcgcgctt 180
ctcgttgtct cctcctccgg caagctctac agcttctccg ccggtgataa cctggtcagg 240
atccttgatc gatatggaaa acagcatgct gatgatctta aagccctgaa tcttcagtca 300
aaagctctga gctatggttc acacaatgag ttacttgaac ttgtggatag caagcttggtg 360
gaatcaaatg tcggtggtgt aagcgtggac accctcgttc agctggaggg tgccttgaa 420
aatgccctct ctctaactag agctaggaag acagaactaa tgttgaagct tgttgatagc 480
ctcaaagaaa aggagaagct gctgaaagaa gagaatcagg ctttggtggtg ccagaaggag 540
aagaagaatc ttgcgggagc cgaagctgat aatatggaga tgtcacctgg acaaatctcc 600
gacatcaatc ttccggtaac tctcccactg cttaattagc caccgttaga cggggctgat 660
caaattaaaa aatccaaaac atacaactaa ataaataagc tttgttggtt ttcacccttg 720
aaggtgacac gttgtatata tcaatactcc cttggctgag agattgtgtg tttactccta 780
tgtagatat aatgagtaaa ataaaaataa aaagatcttt gtaccttcgt cgagagagaa 840
ttgtagttag tgtgcttggt tgttcttttt ctctcttttg cttcatggcg aagaagccta 900
ccgtctaatt tgtaacggag acgtggccct ctctgccctt ttgtattcgt aattcctttg 960
tattaaaaa 969

<210> 4

<211> 868

<212> DNA

<213> Brassica napus

<400> 4

aacgctcagt atctccggct agtggaaacc ggacctcaag atcaaattag ggcgcaaagc 60
actgttgagg acagaagcca tggggaggaa gaaacttgaa atcaagcgaa ttgagaacaa 120
aagtagccga caagttacct tctctaaacg acgcaacggt ctcacgaga aagctcgtca 180

THIS PAGE RI ANK 1115PTO

gctttccggtt ctctgtgacg catccgtcgc tcttctt^ugtc gtctccgcct ccgggaaaact 240
ctacagcttc tcttccggtg ataacctggt caagatcctt gatcgatatg gaaagcaaca 300
tgatgatgat cttaaagcct tggatcgtca gtcaaaagct ttggactgtg gttcacacca 360
tgagctactg gaacttgttg aaagcaagct tgaggaatca aatgtcgata atgtaagtgt 420
gggttccctg gttcagctgg aggaacacct tgagaacgcc ctctccgtaa caagagctag 480
gaagacagaa ctaatgttga agcttgtcga gaaccttaaa gaaaaggaga agttgctgga 540
agaggagaac catgttttgg ctaggcagat ggagaagagt aatcttgtgc gagccgaagc 600
tgataatatg gatgtctcac caggacaaat ctccgacatc aatcttccgg taacgctccc 660
actgcttaat tagtcacctt taatcggcga ataaataaaa tccaaaacat ataactaaaa 720
caaacaagat gtgtaattat ccccttgtaa aggggtgtacg ttgtataatc tatactctct 780
ctccggctcg agaggcttcg ggtgtaaaac tatttcagat ttatgtaaga tagaaaatct 840
atgcaagaca ctttcaaact taaaaaaa 868

<210> 5

<211> 792

<212> DNA

<213> Brassica napus

<400> 5

caaaggcttc tcggagacag aagccatggg aagaaagaaa ctagagatca agcgaattga 60
gaacaaaagt agccgacaag tcaccttctc caaacgacgc aatgggtctca tcgagaaaagc 120
tcgtcagctt tcagttctct gcgatgcac cgctcgtctt ctcgttgtct cagcctccgg 180
caagctttac aacttctccg ccggcgataa cctgggtcaag atccttgatc gatatggaaa 240
acaacatgct gatgatctta aagctctgga tcttcagtca aaagctccga agtatgggtc 300
acaccatgag ctactagagc ttgtcgaag taagcttgtg gaatcaaatt ctgatgtaag 360
cgtcgactcc ctcgttcagc tggaggacca ccttgagact gccctctccg taactagagc 420
taggaagaca gaactaatgt tgaagcttgt tgatagcctc aaagaaaagg agaaattgct 480
gaaagaagag aaccaggggt tggctagcca gatggagaag aataatcttg cgggagccga 540
agctgataaa atggagatgt cacctggaca aatctctgac atcaatcgtc cggtaactct 600
ccgactgctt tattagccnc cttaagtcca aaacttgtga ctaaaaacaa aaataagtta 660
tcgaactatt cccctataag ggtgaacgtt gtatatcttc attctctctg gctgagagac 720
cccgtgtgta aaactatggt tagatttaag taaaaatata tatttaagac atactaaaaa 780
aaaaaaaaaa aa 792

<210> 6

<211> 990

<212> DNA

<213> Brassica napus

<400> 6

gggcacagag accacttgga gacagaaacc atgggaagaa aaaaactaga aatcaagcga 60
attgagaaca aaagtagccg acaagtcacc ttctccaaac gacgcagcgg tctcattgag 120
aaagctcgtc agctttctgt tctctgcgat gcacccgtcg cgcttctcgt tgtctcctcc 180
tccggcaagc tctacagctt ctccgccggt gataacctgg tcaggatcct tgatcgatat 240
ggaaaacagc atgctgatga tcttaaagcc ctgaatcttc agtcaaaagc tctgagctat 300
ggttcacaca atgagttact tgaacttgtg gatagcaagc ttgtggaatc aatgtcgggt 360

THIS PAGE RI ANK (11SDT0)

ggtgtaagcg tggacaccct cgttcagctg gagggtgtcc ttgaaaatgc cctctctcta 420
actagagcta ggaagacaga actaatgttg aagcttggtg atagcctcaa agaaaaggag 480
aagctgctga aagaagagaa tcagggtttg gctggccaga aggagaagaa gaatcttgcg 540
ggagccgaag ctgataatat ggagatgtca cctggacaaa tctccgacat caatcttccg 600
gtaactctcc cactgcttaa ttagccaccg ttagacgggg ctgatcaaat taaaaaatcc 660
aaaacataca actaaataaa taagctttgt tgtttttcac ccttgaaggg tgcacgttgt 720
atatctcaat actcccttgg ctgagagatt gtgtgtttac tcctatgtta gatataatga 780
gtaaaataaa aataaaaaga tctttgtacc ttcgtcgaga gagaattgta gtgagtgtgc 840
ttgtgtgttc tttttctctt ctttgcttca tggcgaagaa gcctaccgtc taatttgtaa 900
cggagacgtg gccctctctg cctttttgta ttcgtaattc ctttgtattt atccacaacg 960
catagagggt gtcattggtt aaaaaaaaaa 990

<210> 7

<211> 780

<212> DNA

<213> Brassica napus

<400> 7

ttagggcaca aaggcttctc ggagacagaa gccatgggaa gaaagaaact agagatcaag 60
cgaattgaga acaaaagtag ccgacaagtc accttctcca aacgacgcaa tgggtctcatc 120
gagaaagctc gtcagctttc agttctctgc gatgcatccg tcgctcttct cgttgtctca 180
gcctccggca agcttttaca cttctccgcc ggcgataacc tgggtcaagat ccttgatcga 240
tatggaaaac aacatgctga tgatcttaaa gctctggatc ttcagtcaaa agctccgaag 300
tatggttcac accatgagct actagagctt gtcgaaagta agcttgtgga atcaaattct 360
gatgtaagcg tcgactccct cgttcagctg gaggaccacc ttgagactgc cctctccgta 420
actagagcta ggaagacaga actaatgttg aagcttggtg atagcctcaa agaaaaggag 480
aaattgctga aagaagagaa ccagggtttg gctagccaga tggagaagaa taatcttgcg 540
ggagccgaag ctgataaaat ggagatgtca cctggacaaa tctctgacat caatcgctccg 600
gtaactctcc gactgcttta ttagccaccct taagtccaaa acttgtgact aaaaacaaaa 660
ataagttatc gaactattcc cctataaggg tgaacgttgt atatcttcat tctctctggc 720
tgagagaccc ccgtgtgtaa actatggnta gatttaagta aaatatatnt ttaagacana 780

<210> 8

<211> 845

<212> DNA

<213> Brassica napus

<400> 8

ctccggctag tggaaaccgg acctcaagat caaattaggg cgcaaagcac tgttggagac 60
agaagccatg gggaggaaga aacttgaaat caagcgaatt gagaacaaaa gtagccgaca 120
agttaccttc tctaaacgac gcaacggtct catcgagaaa gctcgtcagc tttccgttct 180
ctgtgacgca tccgtcgctc ttcttgtcgt ctccgcctcc gggaaactct acagcttctc 240
ctccggtgat aacctggtca agatccttga tcgatatgga aagcaacatg atgatgatct 300
taaagccttg gatcgtcagt caaaagcttt ggactgtggg tcacaccatg agctactgga 360
acttgtggaa agcaagcttg aggaatcaaa tgtcgataat gtaagtgtgg gttccctggg 420
tcagctggag gaacaccttg agaacgccct ctccgtaaca agagctagga agacagaact 480

THIS PAGE RI ANK (ISCPTO)

aatgttgaag cttgtcgaga accttaaaga aaaggagaag ttgctggaag aggagaacca 540
tgttttggct agccagatgg agaagagtaa tcttgtgcga gccgaagctg ataatatgga 600
tgtctcacca ggacaaatct cgcacatcaa tcttccggta acgctccac tgcttaatta 660
gtcaccttta atcggcgaat aaataaaatc caaaacatat aactaaaaca aacaagatgt 720
gtaattatcc ccttgtaaag ggtgtacgtt gtataatcta tactctctct cgggctcgag 780
aggcttcggg tgtaaaacta tttcagattt atgtaagata gaaaatctat gcaagacact 840
ttcaa 845

<210> 9

<211> 825

<212> DNA

<213> Brassica napus

<400> 9

cggcgagagt tgaaaccgaa tctcaggatc aaattagggc acaaaggctt ctccgagaca 60
gaagccatgg gaagaaagaa actagagatc aagcgaattg agaacaaaag tagccgacaa 120
gtcaccttct ccaaaccgacg caatggtctc atcgagaaag ctcgtcagct ttcatgtctc 180
tgcatgcat ccgtcgctct tctcgttgtc tcagcctccg gcaagcttta caacttctcc 240
gccggcgata acctggtcaa gatccttgat cgatatggaa aacaacatgc tgatgatctt 300
aaagctctgg atcttcagtc aaaagctccg aagtatgggt cacaccatga gctactagag 360
cttgtcgaaa gtaagcttgt ggaatcaaat tctgatgtaa gcgtcgactc cctcgttcag 420
ctggaggacc accttgagac tgccctctcc gtaactagag ctaggaagac agaactaatg 480
ttgaagcttg ttgatagcct caaagaaaag gagaaattgc tgaaagaaga gaaccagggt 540
ttggctagcc agatggagaa gaataatctt gcgggagccg aagctgataa aatggagatg 600
tcacctggac aaatctctga catcaatcgt ccggttaact tccgactgct ttattagcca 660
ccttaagtcc aaaacttgtg actaaaaaca aaaataagtt atcgaactat tcccctataa 720
gggtgaacgt tgtatatctt cattctctct ggctgagaga ccccggtgtg aaaactatgg 780
ttagatttaa gtaaaaatat atatttaaga cataactaaa aaaaa 825

<210> 10

<211> 891

<212> DNA

<213> Brassica napus

<400> 10

tccggctagt ggaaaccgga cctcaagatc aaattagggc gcaaagcact gttggagaca 60
gaagccatgg ggaggaagaa acttgaaatc aagcgaattg agaacaaaag tagccgacaa 120
gttaccttct ctaaaccgacg caacggtctc atcgagaaag ctcgtcagct ttccgttctc 180
tgtgacgcat ccgtcgctct tcttgcgtc tccgctccg ggaaactcta cagcttctcc 240
tccggtgata acctggtcaa gatccttgat cgatatggaa agcaacatga tgatgatctt 300
aaagccttgg atcgtcagtc aaaagctttg gactgtgggt cacaccatga gctactggaa 360
cttgtggaaa gcaagcttga ggaatcaaat gtcgataatg taagtgtggg ttccctgggt 420
cagctggagg aacaccttga gaacgcctc tccgtaacaa gagctaggaa gacagaacta 480
atgttgaagc ttgtcgagaa ccttaaagaa aaggagaagt tgctggaaga ggagaacct 540
gttttggcta gccagatgga gaagagtaat cttgtgcgag ccgaagctga taatatggat 600
gtctcaccag gacaaatctc cgacatcaat cttccggtaa cgctccact gcttaattag 660

THIS PAGE RI ANK 111000

tcacctttaa tcggcgaata aataaaatcc a'aaacatata actaaaacaa acaagatgtg 720
 taattatccc cttgtaaagg gtgtacgttg tataatctat actctctctc cggctcgaga 780
 ggcttcgggt gtaaaactat ttcagattta tgtaagatag aaaatctatg caagacactt 840
 tcaaactttg taccttgctt tgtcgacaga gaattacttc gagctaaaaa a 891

<210> 11

<211> 196

<212> PRT

<213> Brassica napus

<400> 11

Met Gly Arg Lys Lys Leu Glu Ile Lys Arg Ile Glu Asn Lys Ser Ser
 1 5 10 15

Arg Gln Val Thr Phe Ser Lys Arg Arg Asn Gly Leu Ile Glu Lys Ala
 20 25 30

Arg Gln Leu Ser Val Leu Cys Asp Ala Ser Val Ala Leu Leu Val Val
 35 40 45

Ser Ala Ser Gly Lys Leu Tyr Asn Phe Ser Ala Gly Asp Asp Leu Val
 50 55 60

Lys Ile Val Asp Arg Tyr Gly Lys Gln His Ala Asp Asp Arg Lys Ala
 65 70 75 80

Leu Asp Leu Gln Ser Glu Ala Pro Lys Tyr Gly Ser His His Glu Leu
 85 90 95

Leu Glu Leu Val Glu Ser Lys Leu Val Glu Ser Asn Ser Asp Val Ser
 100 105 110

Val Asp Ser Leu Val Gln Leu Glu Asn His Leu Glu Thr Ala Leu Ser
 115 120 125

Val Thr Arg Ala Arg Lys Thr Glu Leu Leu Leu Lys Leu Val Asp Ser
 130 135 140

Leu Lys Glu Lys Glu Lys Leu Leu Lys Glu Glu Asn Gln Gly Leu Ala
 145 150 155 160

Ser Gln Met Glu Lys Asn Asn Leu Ala Gly Ala Glu Ala Asp Lys Met
 165 170 175

Glu Val Ser Pro Gly Gln Ile Ser Asp Ile Asn Cys Pro Val Thr Leu
 180 185 190

THIS PAGE RI ANK (15PTO)

Pro Leu Leu Tyr
195

<210> 12
<211> 196
<212> PRT
<213> Brassica napus

<400> 12

Met Gly Arg Lys Lys Leu Glu Ile Lys Arg Ile Glu Lys Asn Ser Ser
1 5 10 15

Arg Gln Val Thr Phe Cys Lys Arg Arg Asn Gly Leu Ile Glu Lys Ala
20 25 30

Arg Gln Leu Ser Val Leu Cys Glu Ala Ser Val Gly Leu Leu Val Val
35 40 45

Ser Ala Ser Asp Lys Leu Tyr Ser Phe Ser Ser Gly Asp Arg Leu Glu
50 55 60

Lys Ile Leu Asp Arg Tyr Gly Lys Lys His Ala Asp Asp Leu Asn Ala
65 70 75 80

Leu Asp Leu Gln Ser Lys Ser Leu Asn Tyr Ser Ser His His Glu Leu
85 90 95

Leu Glu Leu Val Glu Ser Lys Leu Val Glu Ser Ile Asp Asp Val Ser
100 105 110

Val Asp Ser Leu Val Glu Leu Glu Asp His Leu Glu Thr Ala Leu Ser
115 120 125

Val Thr Arg Ala Arg Lys Ala Glu Leu Met Leu Lys Leu Val Glu Ser
130 135 140

Leu Lys Glu Lys Glu Asn Leu Leu Lys Glu Glu Asn Gln Val Leu Ala
145 150 155 160

Ser Gln Ile Glu Lys Lys Asn Leu Glu Gly Ala Glu Ala Asp Asn Ile
165 170 175

Glu Met Ser Ser Gly Gln Ile Ser Asp Ile Asn Leu Pro Val Thr Leu
180 185 190

Pro Leu Leu Asn
195

THIS PAGE RI ANK (ISPTO)

<210> 13

<211> 197

<212> PRT

<213> Brassica napus

<400> 13

Met Gly Arg Lys Lys Leu Glu Ile Lys Arg Ile Glu Asn Lys Ser Ser
1 5 10 15

Arg Gln Val Thr Phe Ser Lys Arg Arg Ser Gly Leu Ile Glu Lys Ala
20 25 30

Arg Gln Leu Ser Val Leu Cys Asp Ala Ser Val Ala Leu Leu Val Val
35 40 45

Ser Ser Ser Gly Lys Leu Tyr Ser Phe Ser Ala Gly Asp Asn Leu Val
50 55 60

Arg Ile Leu Asp Arg Tyr Gly Lys Gln His Ala Asp Asp Leu Lys Ala
65 70 75 80

Leu Asn Leu Gln Ser Lys Ala Leu Ser Tyr Gly Ser His Asn Glu Leu
85 90 95

Leu Glu Leu Val Asp Ser Lys Leu Val Glu Ser Asn Val Gly Gly Val
100 105 110

Ser Val Asp Thr Leu Val Gln Leu Glu Gly Val Leu Glu Asn Ala Leu
115 120 125

Ser Leu Thr Arg Ala Arg Lys Thr Glu Leu Met Leu Lys Leu Val Asp
130 135 140

Ser Leu Lys Glu Lys Glu Lys Leu Leu Lys Glu Glu Asn Gln Ala Leu
145 150 155 160

Ala Gly Gln Lys Glu Lys Lys Asn Leu Ala Gly Ala Glu Ala Asp Asn
165 170 175

Met Glu Met Ser Pro Gly Gln Ile Ser Asp Ile Asn Leu Pro Val Thr
180 185 190

Leu Pro Leu Leu Asn
195

THIS PAGE BLANK (USPTO)

<210> 14
<211> 197
<212> PRT
<213> Brassica napus

<400> 14

Met Gly Arg Lys Lys Leu Glu Ile Lys Arg Ile Glu Asn Lys Ser Ser
1 5 10 15

Arg Gln Val Thr Phe Ser Lys Arg Arg Asn Gly Leu Ile Glu Lys Ala
20 25 30

Arg Gln Leu Ser Val Leu Cys Asp Ala Ser Val Ala Leu Leu Val Val
35 40 45

Ser Ala Ser Gly Lys Leu Tyr Ser Phe Ser Ser Gly Asp Asn Leu Val
50 55 60

Lys Ile Leu Asp Arg Tyr Gly Lys Gln His Asp Asp Asp Leu Lys Ala
65 70 75 80

Leu Asp Arg Gln Ser Lys Ala Leu Asp Cys Gly Ser His His Glu Leu
85 90 95

Leu Glu Leu Val Glu Ser Lys Leu Glu Glu Ser Asn Val Asp Asn Val
100 105 110

Ser Val Gly Ser Leu Val Gln Leu Glu Glu His Leu Glu Asn Ala Leu
115 120 125

Ser Val Thr Arg Ala Arg Lys Thr Glu Leu Met Leu Lys Leu Val Glu
130 135 140

Asn Leu Lys Glu Lys Glu Lys Leu Leu Glu Glu Glu Asn His Val Leu
145 150 155 160

Ala Ser Gln Met Glu Lys Ser Asn Leu Val Arg Ala Glu Ala Asp Asn
165 170 175

Met Asp Val Ser Pro Gly Gln Ile Ser Asp Ile Asn Leu Pro Val Thr
180 185 190

Leu Pro Leu Leu Asn
195

<210> 15
<211> 196

THIS PAGE BLANK (USPTO)

<212> PRT

<213> Brassica napus

<400> 15

Met Gly Arg Lys Lys Leu Glu Ile Lys Arg Ile Glu Asn Lys Ser Ser
1 5 10 15

Arg Gln Val Thr Phe Ser Lys Arg Arg Asn Gly Leu Ile Glu Lys Ala
20 25 30

Arg Gln Leu Ser Val Leu Cys Asp Ala Ser Val Ala Leu Leu Val Val
35 40 45

Ser Ala Ser Gly Lys Leu Tyr Asn Phe Ser Ala Gly Asp Asn Leu Val
50 55 60

Lys Ile Leu Asp Arg Tyr Gly Lys Gln His Ala Asp Asp Leu Lys Ala
65 70 75 80

Leu Asp Leu Gln Ser Lys Ala Pro Lys Tyr Gly Ser His His Glu Leu
85 90 95

Leu Glu Leu Val Glu Ser Lys Leu Val Glu Ser Asn Ser Asp Val Ser
100 105 110

Val Asp Ser Leu Val Gln Leu Glu Asp His Leu Glu Thr Ala Leu Ser
115 120 125

Val Thr Arg Ala Arg Lys Thr Glu Leu Met Leu Lys Leu Val Asp Ser
130 135 140

Leu Lys Glu Lys Glu Lys Leu Leu Lys Glu Glu Asn Gln Gly Leu Ala
145 150 155 160

Ser Gln Met Glu Lys Asn Asn Leu Ala Gly Ala Glu Ala Asp Lys Met
165 170 175

Glu Met Ser Pro Gly Gln Ile Ser Asp Ile Asn Arg Pro Val Thr Leu
180 185 190

Arg Leu Leu Tyr
195

<210> 16

<211> 197

<212> PRT

<213> Brassica napus

THIS PAGE BLANK (DISPTD)

<400> 16

Met Gly Arg Lys Lys Leu Glu Ile Lys Arg Ile Glu Asn Lys Ser Ser
1 5 10 15

Arg Gln Val Thr Phe Ser Lys Arg Arg Ser Gly Leu Ile Glu Lys Ala
20 25 30

Arg Gln Leu Ser Val Leu Cys Asp Ala Ser Val Ala Leu Leu Val Val
35 40 45

Ser Ser Ser Gly Lys Leu Tyr Ser Phe Ser Ala Gly Asp Asn Leu Val
50 55 60

Arg Ile Leu Asp Arg Tyr Gly Lys Gln His Ala Asp Asp Leu Lys Ala
65 70 75 80

Leu Asn Leu Gln Ser Lys Ala Leu Ser Tyr Gly Ser His Asn Glu Leu
85 90 95

Leu Glu Leu Val Asp Ser Lys Leu Val Glu Ser Asn Val Gly Gly Val
100 105 110

Ser Val Asp Thr Leu Val Gln Leu Glu Gly Val Leu Glu Asn Ala Leu
115 120 125

Ser Leu Thr Arg Ala Arg Lys Thr Glu Leu Met Leu Lys Leu Val Asp
130 135 140

Ser Leu Lys Glu Lys Glu Lys Leu Leu Lys Glu Glu Asn Gln Ala Leu
145 150 155 160

Ala Gly Gln Lys Glu Lys Lys Asn Leu Ala Gly Ala Glu Ala Asp Asn
165 170 175

Met Glu Met Ser Pro Gly Gln Ile Ser Asp Ile Asn Leu Pro Val Thr
180 185 190

Leu Pro Leu Leu Asn
195

<210> 17

<211> 196

<212> PRT

<213> Brassica napus

<400> 17

THIS PAGE RI ANK (ISPTO)

Met Gly Arg Lys Lys Leu Glu Ile Lys Arg Ile Glu Asn Lys Ser Ser
 1 5 10 15

Arg Gln Val Thr Phe Ser Lys Arg Arg Asn Gly Leu Ile Glu Lys Ala
 20 25 30

Arg Gln Leu Ser Val Leu Cys Asp Ala Ser Val Ala Leu Leu Val Val
 35 40 45

Ser Ala Ser Gly Lys Leu Tyr Asn Phe Ser Ala Gly Asp Asn Leu Val
 50 55 60

Lys Ile Leu Asp Arg Tyr Gly Lys Gln His Ala Asp Asp Leu Lys Ala
 65 70 75 80

Leu Asp Leu Gln Ser Lys Ala Pro Lys Tyr Gly Ser His His Glu Leu
 85 90 95

Leu Glu Leu Val Glu Ser Lys Leu Val Glu Ser Asn Ser Asp Val Ser
 100 105 110

Val Asp Ser Leu Val Gln Leu Glu Asp His Leu Glu Thr Ala Leu Ser
 115 120 125

Val Thr Arg Ala Arg Lys Thr Glu Leu Met Leu Lys Leu Val Asp Ser
 130 135 140

Leu Lys Glu Lys Glu Lys Leu Leu Lys Glu Glu Asn Gln Gly Leu Ala
 145 150 155 160

Ser Gln Met Glu Lys Asn Asn Leu Ala Gly Ala Glu Ala Asp Lys Met
 165 170 175

Glu Met Ser Pro Gly Gln Ile Ser Asp Ile Asn Arg Pro Val Thr Leu
 180 185 190

Arg Leu Leu Tyr
 195

<210> 18

<211> 197

<212> PRT

<213> Brassica napus

<400> 18

Met Gly Arg Lys Lys Leu Glu Ile Lys Arg Ile Glu Asn Lys Ser Ser
 1 5 10 15

THIS PAGE RI ANK 1115PTO

Arg Gln Val Thr Phe Ser Lys Arg Arg Asn Gly Leu Ile Glu Lys Ala
 20 25 30

Arg Gln Leu Ser Val Leu Cys Asp Ala Ser Val Ala Leu Leu Val Val
 35 40 45

Ser Ala Ser Gly Lys Leu Tyr Ser Phe Ser Ser Gly Asp Asn Leu Val
 50 55 60

Lys Ile Leu Asp Arg Tyr Gly Lys Gln His Asp Asp Asp Leu Lys Ala
 65 70 75 80

Leu Asp Arg Gln Ser Lys Ala Leu Asp Cys Gly Ser His His Glu Leu
 85 90 95

Leu Glu Leu Val Glu Ser Lys Leu Glu Glu Ser Asn Val Asp Asn Val
 100 105 110

Ser Val Gly Ser Leu Val Gln Leu Glu Glu His Leu Glu Asn Ala Leu
 115 120 125

Ser Val Thr Arg Ala Arg Lys Thr Glu Leu Met Leu Lys Leu Val Glu
 130 135 140

Asn Leu Lys Glu Lys Glu Lys Leu Leu Glu Glu Glu Asn His Val Leu
 145 150 155 160

Ala Ser Gln Met Glu Lys Ser Asn Leu Val Arg Ala Glu Ala Asp Asn
 165 170 175

Met Asp Val Ser Pro Gly Gln Ile Ser Asp Ile Asn Leu Pro Val Thr
 180 185 190

Leu Pro Leu Leu Asn
 195

<210> 19

<211> 196

<212> PRT

<213> Brassica napus

<400> 19

Met Gly Arg Lys Lys Leu Glu Ile Lys Arg Ile Glu Asn Lys Ser Ser
 1 5 10 15

Arg Gln Val Thr Phe Ser Lys Arg Arg Asn Gly Leu Ile Glu Lys Ala

THIS PAGE RI ANK (115PTO)

20

25

30

Arg Gln Leu Ser Val Leu Cys Asp Ala Ser Val Ala Leu Leu Val Val
 35 40 45

Ser Ala Ser Gly Lys Leu Tyr Asn Phe Ser Ala Gly Asp Asn Leu Val
 50 55 60

Lys Ile Leu Asp Arg Tyr Gly Lys Gln His Ala Asp Asp Leu Lys Ala
 65 70 75 80

Leu Asp Leu Gln Ser Lys Ala Pro Lys Tyr Gly Ser His His Glu Leu
 85 90 95

Leu Glu Leu Val Glu Ser Lys Leu Val Glu Ser Asn Ser Asp Val Ser
 100 105 110

Val Asp Ser Leu Val Gln Leu Glu Asp His Leu Glu Thr Ala Leu Ser
 115 120 125

Val Thr Arg Ala Arg Lys Thr Glu Leu Met Leu Lys Leu Val Asp Ser
 130 135 140

Leu Lys Glu Lys Glu Lys Leu Leu Lys Glu Glu Asn Gln Gly Leu Ala
 145 150 155 160

Ser Gln Met Glu Lys Asn Asn Leu Ala Gly Ala Glu Ala Asp Lys Met
 165 170 175

Glu Met Ser Pro Gly Gln Ile Ser Asp Ile Asn Arg Pro Val Thr Leu
 180 185 190

Arg Leu Leu Tyr
 195

<210> 20

<211> 197

<212> PRT

<213> Brassica napus

<400> 20

Met Gly Arg Lys Lys Leu Glu Ile Lys Arg Ile Glu Asn Lys Ser Ser
 1 5 10 15

Arg Gln Val Thr Phe Ser Lys Arg Arg Asn Gly Leu Ile Glu Lys Ala
 20 25 30

THIS PAGE BLANK (11/15/70)

Arg Gln Leu Ser Val Leu Cys Asp Ala Ser Val Ala Leu Leu Val Val
 35 40 45

Ser Ala Ser Gly Lys Leu Tyr Ser Phe Ser Ser Gly Asp Asn Leu Val
 50 55 60

Lys Ile Leu Asp Arg Tyr Gly Lys Gln His Asp Asp Asp Leu Lys Ala
 65 70 75 80

Leu Asp Arg Gln Ser Lys Ala Leu Asp Cys Gly Ser His His Glu Leu
 85 90 95

Leu Glu Leu Val Glu Ser Lys Leu Glu Glu Ser Asn Val Asp Asn Val
 100 105 110

Ser Val Gly Ser Leu Val Gln Leu Glu Glu His Leu Glu Asn Ala Leu
 115 120 125

Ser Val Thr Arg Ala Arg Lys Thr Glu Leu Met Leu Lys Leu Val Glu
 130 135 140

Asn Leu Lys Glu Lys Glu Lys Leu Leu Glu Glu Glu Asn His Val Leu
 145 150 155 160

Ala Ser Gln Met Glu Lys Ser Asn Leu Val Arg Ala Glu Ala Asp Asn
 165 170 175

Met Asp Val Ser Pro Gly Gln Ile Ser Asp Ile Asn Leu Pro Val Thr
 180 185 190

Leu Pro Leu Leu Asn
 195

<210> 21

<211> 196

<212> PRT

<213> Brassica napus

<400> 21

Met Gly Arg Arg Lys Ile Glu Ile Lys Arg Ile Glu Asn Lys Ser Ser
 1 5 10 15

Arg Gln Val Thr Phe Ser Lys Arg Arg Asn Gly Leu Ile Asp Lys Ala
 20 25 30

Arg Gln Leu Ser Ile Leu Cys Glu Ser Ser Val Ala Val Val Val Val
 35 40 45

THIS PAGE RI ANK (15PTO)

Ser Ala Ser Gly Lys Leu Tyr Asp Ser Ser Ser Gly Asp Asp Ile Ser
50 55 60

Lys Ile Ile Asp Arg Tyr Glu Ile Gln His Ala Asp Glu Leu Arg Ala
65 70 75 80

Leu Asp Leu Glu Glu Lys Ile Gln Asn Tyr Leu Pro His Lys Glu Leu
85 90 95

Leu Glu Thr Val Gln Ser Lys Leu Glu Glu Pro Asn Val Asp Asn Val
100 105 110

Ser Val Asp Ser Leu Ile Ser Leu Glu Glu Gln Leu Glu Thr Ala Leu
115 120 125

Ser Val Ser Arg Ala Arg Lys Ala Glu Leu Met Met Glu Tyr Ile Glu
130 135 140

Ser Leu Lys Glu Lys Glu Lys Leu Leu Arg Glu Glu Asn Gln Val Leu
145 150 155 160

Ala Ser Gln Met Gly Lys Asn Thr Leu Leu Ala Thr Asp Asp Glu Arg
165 170 175

Gly Met Phe Pro Gly Ser Ser Ser Gly Asn Lys Ile Pro Glu Thr Leu
180 185 190

Pro Leu Leu Asn
195

<210> 22

<211> 196

<212> PRT

<213> Brassica napus

<400> 22

Met Gly Arg Arg Lys Val Glu Ile Lys Arg Ile Glu Asn Lys Ser Ser
1 5 10 15

Arg Gln Val Thr Phe Ser Lys Arg Arg Lys Gly Leu Ile Glu Lys Ala
20 25 30

Arg Gln Leu Ser Ile Leu Cys Glu Ser Ser Ile Ala Val Val Ala Val
35 40 45

Ser Gly Ser Gly Lys Leu Tyr Asp Ser Ala Ser Gly Asp Asn Met Ser

THIS PAGE RI ANK (115PTO)

50 55 60
 Lys Ile Ile Asp Arg Tyr Glu Ile His His Ala Asp Glu Leu Lys Ala
 65 70 75 80
 Leu Asp Leu Ala Glu Lys Ile Arg Asn Tyr Leu Pro His Lys Glu Leu
 85 90 95
 Leu Glu Ile Val Gln Ser Lys Leu Glu Glu Ser Asn Val Asp Asn Val
 100 105 110
 Ser Val Asp Ser Leu Ile Ser Met Glu Glu Gln Leu Glu Thr Ala Leu
 115 120 125
 Ser Val Ile Arg Ala Lys Lys Thr Glu Leu Met Met Glu Asp Met Lys
 130 135 140
 Ser Leu Gln Glu Arg Glu Lys Leu Leu Ile Glu Glu Asn Gln Ile Leu
 145 150 155 160
 Ala Ser Gln Val Gly Lys Lys Thr Phe Leu Val Ile Glu Gly Asp Arg
 165 170 175
 Gly Met Ser Arg Glu Asn Gly Ser Gly Asn Lys Val Pro Glu Thr Leu
 180 185 190
 Ser Leu Leu Lys
 195

<210> 23

<211> 200

<212> PRT

<213> Brassica napus

<400> 23

Met Gly Arg Arg Lys Val Glu Ile Lys Arg Ile Glu Asn Lys Ser Ser
 1 5 10 15
 Arg Gln Val Thr Phe Cys Lys Arg Arg Asn Gly Leu Met Glu Lys Ala
 20 25 30
 Arg Gln Leu Ser Ile Leu Cys Glu Ser Ser Val Ala Leu Ile Ile Ile
 35 40 45
 Ser Ala Thr Gly Arg Leu Tyr Ser Phe Ser Ser Gly Asp Ser Met Ala
 50 55 60

01-00-00

THIS PAGE BLANK (ISPTO)

Lys Ile Leu Ser Arg Tyr Glu Leu Glu Gln Ala Asp Asp Leu Lys Thr
65 70 75 80

Leu Asp Leu Glu Glu Lys Thr Leu Asn Tyr Leu Ser His Lys Glu Leu
85 90 95

Leu Glu Thr Ile Gln Cys Lys Ile Glu Glu Ala Lys Ser Asp Asn Val
100 105 110

Ser Ile Asp Cys Leu Lys Ser Leu Glu Glu Gln Leu Lys Thr Ala Leu
115 120 125

Ser Val Thr Arg Ala Arg Lys Thr Glu Leu Met Met Glu Leu Val Lys
130 135 140

Thr His Gln Glu Lys Glu Lys Leu Leu Arg Glu Glu Asn Gln Ser Leu
145 150 155 160

Thr Asn Gln Leu Ile Lys Met Gly Lys Met Lys Lys Ser Val Glu Ala
165 170 175

Glu Asp Ala Arg Ala Met Ser Pro Glu Ser Ser Ser Asp Asn Lys Pro
180 185 190

Pro Glu Thr Leu Leu Leu Leu Lys
195 200

<210> 24

<211> 198

<212> PRT

<213> Brassica napus

<400> 24

Met Gly Arg Arg Arg Val Glu Ile Lys Arg Ile Glu Asn Lys Ser Ser
1 5 10 15

Arg Gln Val Thr Phe Cys Lys Arg Arg Asn Gly Leu Met Glu Lys Ala
20 25 30

Arg Gln Leu Ser Ile Leu Cys Gly Ser Ser Val Ala Leu Phe Ile Val
35 40 45

Ser Ser Thr Gly Lys Leu Tyr Asn Ser Ser Ser Gly Asp Ser Met Ala
50 55 60

Lys Ile Ile Ser Arg Phe Lys Ile Gln Gln Ala Asp Asp Pro Glu Thr
65 70 75 80

THIS PAGE BLANK (ISPTO)

Leu Asp Leu Glu Asp Lys Thr Gln Asp Tyr Leu Ser His Lys Glu Leu
85 90 95

Leu Glu Ile Val Gln Arg Lys Ile Glu Glu Ala Lys Gly Asp Asn Val
100 105 110

Ser Ile Glu Ser Leu Ile Ser Met Glu Glu Gln Leu Lys Ser Ala Leu
115 120 125

Ser Val Ile Arg Ala Arg Lys Thr Glu Leu Leu Met Glu Leu Val Lys
130 135 140

Asn Leu Gln Asp Lys Glu Lys Leu Leu Lys Glu Lys Asn Lys Val Leu
145 150 155 160

Ala Ser Glu Val Gly Lys Leu Lys Lys Ile Leu Glu Thr Gly Asp Glu
165 170 175

Arg Ala Val Met Ser Pro Glu Asn Ser Ser Gly His Ser Pro Pro Glu
180 185 190

Thr Leu Pro Leu Leu Lys
195

<210> 25

<211> 196

<212> PRT

<213> Brassica napus

<400> 25

Met Gly Arg Lys Lys Val Glu Ile Lys Arg Ile Glu Asn Lys Ser Ser
1 5 10 15

Arg Gln Val Thr Phe Ser Lys Arg Arg Asn Gly Leu Ile Glu Lys Ala
20 25 30

Arg Gln Leu Ser Ile Leu Cys Glu Ser Ser Ile Ala Val Leu Val Val
35 40 45

Ser Gly Ser Gly Lys Leu Tyr Lys Ser Ala Ser Gly Asp Asn Met Ser
50 55 60

Lys Ile Ile Asp Arg Tyr Glu Ile His His Ala Asp Glu Leu Glu Ala
65 70 75 80

Leu Asp Leu Ala Glu Lys Thr Arg Asn Tyr Leu Pro Leu Lys Glu Leu

THIS PAGE BLANK (ISPTO)

85

. 90^

95

Leu Glu Ile Val Gln Ser Lys Leu Glu Glu Ser Asn Val Asp Asn Ala
 100 105 110

Ser Val Asp Thr Leu Ile Ser Leu Glu Glu Gln Leu Glu Thr Ala Leu
 115 120 125

Ser Val Thr Arg Ala Arg Lys Thr Glu Leu Met Met Gly Glu Val Lys
 130 135 140

Ser Leu Gln Lys Thr Glu Asn Leu Leu Arg Glu Glu Asn Gln Thr Leu
 145 150 155 160

Ala Ser Gln Val Gly Lys Lys Thr Phe Leu Val Ile Glu Gly Asp Arg
 165 170 175

Gly Met Ser Trp Glu Asn Gly Ser Gly Asn Lys Val Arg Glu Thr Leu
 180 185 190

Pro Leu Leu Lys
 195

<210> 26

<211> 691

<212> DNA

<213> Brassica napus

<400> 26

ggatcctatg tgaggttata gatccacgtg catatcaaatt ttctcaaatt cgttgccttc 60
 taacggctct tatatccctt ctccgtggac agaattcttca gtcaaaagct ctgagctatg 120
 gttcacacaa tgagttactt gaacttgttg atagggttagt actacctgag acttttttcc 180
 ctctctctct ttcattacaa aggtattagg gtttcttgtc aatctgtgca tatatatgca 240
 gcaagcttgt ggaatcaaatt gtcggtggtg taagcgtgga caccctcgtt cagctggagg 300
 gtgtccttga aaatgccctc tctctaacta gagctaggaa ggtacgttga cttcatactg 360
 tcttctcatt tcttactttg tttgttgaaa cgattgttca cttatattta atttgttgca 420
 gacagaacta atgttgaagc ttgttgatag cctcaaagaa aagggttagat atatcatata 480
 tgattttata gcacttcaga tatcttctcg tgtttgaaag cctcaaatat ttatgtgttg 540
 tattaagttt ctctaagtgt gctttatgag ctgcgaatca aacttcttca taagtgcac 600
 tgggtctttca nggatgatta aaaatattgt tttggataacc agaactctgaa aatangntta 660
 aaaacttgca actgatgaac atgtccttca n 691

<210> 27

<211> 68

<212> PRT

<213> Brassica napus

THIS PAGE BLANK (USPTO)

<400> 27

Asn Leu Gln Ser Lys Ala Leu Ser Tyr Gly Ser His Asn Glu Leu Leu
1 5 10 15

Glu Leu Val Asp Ser Lys Leu Val Glu Ser Asn Val Gly Gly Val Ser
20 25 30

Val Asp Thr Leu Val Gln Leu Glu Gly Val Leu Glu Asn Ala Leu Ser
35 40 45

Leu Thr Arg Ala Arg Lys Thr Glu Leu Met Leu Lys Leu Val Asp Ser
50 55 60

Leu Lys Glu Lys
65

THIS PAGE BLANK (USPTO)

INTERNATIONAL SEARCH REPORT

International application No.
PCT/AU 99/01079

A. CLASSIFICATION OF SUBJECT MATTER		
Int Cl ⁷ : C12N 15/29		
According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED		
Minimum documentation searched (classification system followed by classification symbols) SEE ELECTRONIC DATABASE BOX BELOW		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched SEE ELECTRONIC DATABASE BOX BELOW		
Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) Medline, Chem Abs, WPID: MADS box, inflorescence, flower, control, development, DNA binding, genes, FLF, flower locus.		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 94/00582 (Centre for Plant Breeding and Reproduction Research) 6 January 1994 see entire document	1-23
X	WO 96/11566 (Washington State University) 25 April 1996 see entire document	1-23
PX	SHELDON, C.C et al (March 1999) "The <i>FLF</i> MADS Box Gene: A Repressor of Flowering in Arabidopsis Regulated by Vernalization and Methylation" <i>The Plant Cell</i> , vol. 11, pages 445-458. see entire document	1-23
<input type="checkbox"/> Further documents are listed in the continuation of Box C <input checked="" type="checkbox"/> See patent family annex		
<p>* Special categories of cited documents:</p> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier application or patent but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p> <p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art</p> <p>"&" document member of the same patent family</p>		
Date of the actual completion of the international search 9 February 2000		Date of mailing of the international search report 11 FEB 2000
Name and mailing address of the ISA/AU AUSTRALIAN PATENT OFFICE PO BOX 200, WODEN ACT 2606, AUSTRALIA E-mail address: pct@ipaustalia.gov.au Facsimile No. (02) 6285 3929		Authorized officer PHILIPPA WYRDEMAN Telephone No.: (02) 6283 2554

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No.
PCT/AU 99/01079

This Annex lists the known "A" publication level patent family members relating to the patent documents cited in the above-mentioned international search report. The Australian Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

Patent Document Cited in Search Report	Patent Family Member
WO 96/11566 AU 39569/95 EP 784425	
WO 94/00582 EP 672155	
END OF ANNEX	

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

☐ BLACK BORDERS

☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES

☐ FADED TEXT OR DRAWING

☒ BLURRED OR ILLEGIBLE TEXT OR DRAWING

☐ SKEWED/SLANTED IMAGES

☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS

☐ GRAY SCALE DOCUMENTS

☒ LINES OR MARKS ON ORIGINAL DOCUMENT

☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY

☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.

THIS PAGE BLANK (USPTO)